

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 237/32, C07D 241/24, 213/82, A61K 31/495, 31/44	A1	(11) International Publication Number: WO 99/50230	
		(43) International Publication Date: 7 October 1999 (07.10.99)	
(21) International Application Number: PCT/US99/07149 (22) International Filing Date: 31 March 1999 (31.03.99) (30) Priority Data: 60/080,060 31 March 1998 (31.03.98) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 60/080,060 (CON) Filed on 31 March 1998 (31.03.98) (71) Applicant (for all designated States except US): VERTEX PHARMACEUTICALS INCORPORATED [US/US]; 130 Waverly Street, Cambridge, MA 02139-4242 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): TUNG, Roger, D. [US/US]; 54 Richfield Road, Arlington, MA 01274 (US). FARMER, Luc, J. [CA/US]; 19 Howe Lane, Foxboro, MA 02035 (US). BHISETTI, Govinda, R. [IN/US]; 70 Grassland Street, Lexington, MA 02173 (US). (74) Agents: HALEY, James, F., Jr.; Fish & Neave, 1251 Avenue of the Americas, New York, NY 10020 (US) et al.		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: INHIBITORS OF SERINE PROTEASES, PARTICULARLY HEPATITIS C VIRUS NS3 PROTEASE			
(57) Abstract The present invention relates to compounds, methods and pharmaceutical compositions for inhibiting proteases, particularly serine proteases, and more particularly HCV NS3 proteases. The compounds, and the compositions and methods that utilize them, can be used, either alone or in combination to inhibit viruses, particularly HCV virus.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INHIBITORS OF SERINE PROTEASES,
PARTICULARLY HEPATITIS C VIRUS NS3 PROTEASE

TECHNICAL FIELD OF THE INVENTION

5 The present invention relates to compounds that are useful as protease inhibitors, particularly as serine protease inhibitors, and more particularly as hepatitis C NS3 protease inhibitors. As such, they act by interfering with the life cycle of the hepatitis C virus
10 and are also useful as antiviral agents.

 This invention also relates to pharmaceutical compositions comprising these compounds. The compounds and pharmaceutical compositions of this invention are particularly well suited for inhibiting HCV NS3 protease
15 activity and consequently, may be advantageously used as therapeutic agents against the hepatitis C virus and other viruses that are dependent upon a serine protease for proliferation. This invention also relates to methods for inhibiting the activity of proteases,
20 including hepatitis C virus NS3 protease and other serine proteases, using the compounds of this invention and related compounds.

BACKGROUND OF THE INVENTION

 Infection by hepatitis C virus ("HCV") is a
25 compelling human medical problem. HCV is recognized as the causative agent for most cases of non-A, non-B hepatitis, with an estimated human seroprevalence of 1%

globally [Purcell, R.H., "Hepatitis C virus: Historical perspective and current concepts" FEMS Microbiology Reviews 14, pp. 181-192 (1994); Van der Poel, C.L., "Hepatitis C Virus. Epidemiology, Transmission and Prevention in Hepatitis C Virus. Current Studies in Hematology and Blood Transfusion, H.W. Reesink, Ed., (Basel: Karger), pp. 137-163 (1994)]. Four million individuals may be infected in the United States alone [Alter, M.J. and Mast, E.E., "The Epidemiology of Viral Hepatitis in the United States, Gastroenterol. Clin. North Am. 23, pp. 437-455 (1994)].

Upon first exposure to HCV only about 20% of infected individuals develop acute clinical hepatitis while others appear to resolve the infection spontaneously. In most instances, however, the virus establishes a chronic infection that persists for decades [Iwarson, S. "The Natural Course of Chronic Hepatitis" FEMS Microbiology Reviews 14, pp. 201-204 (1994)]. This usually results in recurrent and progressively worsening liver inflammation, which often leads to more severe disease states such as cirrhosis and hepatocellular carcinoma [Kew, M.C., "Hepatitis C and Hepatocellular Carcinoma", FEMS Microbiology Reviews, 14, pp. 211-220 (1994); Saito, I., et al. "Hepatitis C Virus Infection is Associated with the Development of Hepatocellular Carcinoma" Proc. Natl. Acad. Sci. USA 87, pp. 6547-6549 (1990)]. Unfortunately, there are no broadly effective treatments for the debilitating progression of chronic HCV.

The HCV genome encodes a polyprotein of 3010-3033 amino acids [Choo, Q.-L., et al. "Genetic Organization and Diversity of the Hepatitis C Virus",

Proc. Natl. Acad. Sci. USA, 88, pp. 2451-2455 (1991); Kato, N. et al., Molecular Cloning of the Human Hepatitis C Virus Genome From Japanese Patients with Non-A, Non-B Hepatitis", Proc. Natl. Acad. Sci. USA, 87, pp. 9524-9528
5 (1990); Takamizawa, A. et al., "Structure and Organization of the Hepatitis C Virus Genome Isolated From Human Carriers", J. Virol., 65, pp. 1105-1113 (1991)]. The HCV nonstructural (NS) proteins are presumed to provide the essential catalytic machinery for
10 viral replication. The NS proteins are derived by proteolytic cleavage of the polyprotein [Bartenschlager, R. et al., "Nonstructural Protein 3 of the Hepatitis C Virus Encodes a Serine-Type Proteinase Required for Cleavage at the NS3/4 and NS4/5 Junctions", J. Virol.,
15 67, pp. 3835-3844 (1993); Grakoui, A. et al. "Characterization of the Hepatitis C Virus-Encoded Serine Proteinase: Determination of Proteinase-Dependent Polyprotein Cleavage Sites", J. Virol., 67, pp. 2832-2843 (1993); Grakoui, A. et al., Expression and Identification
20 of Hepatitis C Virus Polyprotein Cleavage Products", J. Virol., 67, pp. 1385-1395 (1993); Tomei, L. et al., "NS3 is a serine protease required for processing of hepatitis C virus polyprotein", J. Virol., 67, pp. 4017-4026 (1993)].

25 The HCV NS protein 3 (NS3) contains a serine protease activity that helps process the majority of the viral enzymes, and is thus considered essential for viral replication and infectivity. It is known that mutations in the yellow fever virus NS3 protease decreases viral
30 infectivity [Chambers, T.J. et. al., "Evidence that the N-terminal Domain of Nonstructural Protein NS3 From Yellow Fever Virus is a Serine Protease Responsible for

Site-Specific Cleavages in the Viral Polyprotein", Proc. Natl. Acad. Sci. USA, 87, pp. 8898-8902 (1990)]. The first 181 amino acids of NS3 (residues 1027-1207 of the viral polyprotein) have been shown to contain the serine protease domain of NS3 that processes all four downstream sites of the HCV polyprotein [C. Lin et al., "Hepatitis C Virus NS3 Serine Proteinase: Trans-Cleavage Requirements and Processing Kinetics", J. Virol., 68, pp. 8147-8157 (1994)].

10 The HCV NS3 serine protease and its associated cofactor, NS4A, helps process all of the viral enzymes, and is thus considered essential for viral replication. This processing appears to be analogous to that carried out by the human immunodeficiency virus aspartyl
15 protease, which is also involved in viral enzyme processing HIV protease inhibitors, which inhibit viral protein processing are potent antiviral agents in man, indicating that interrupting this stage of the viral life cycle results in therapeutically active agents.
20 Consequently it is an attractive target for drug discovery. Unfortunately, there are no serine protease inhibitors available currently as anti-HCV agents.

 Furthermore, the current understanding of HCV has not led to any other satisfactory anti-HCV agents or
25 treatments. The only established therapy for HCV disease is interferon treatment. However, interferons have significant side effects (Janssen et al., 1994; Renault and Hoofnagle, 1989) [Janssen, H. L. A., et al. "Suicide Associated with Alfa-Interferon Therapy for Chronic Viral
30 Hepatitis" J. Hepatol., 21, pp. 241-243 (1994)]; Renault, P.F. and Hoofnagle, J.H., "Side effects of alpha interferon. Seminars in Liver Disease 9, 273-277. (1989)]

and induce long term remission in only a fraction (~ 25%) of cases [Weiland, O. "Interferon Therapy in Chronic Hepatitis C Virus Infection", FEMS Microbiol. Rev., 14, PP. 279-288 (1994)]. Moreover, the prospects for effective anti-HCV vaccines remain uncertain.

Thus, there is a need for more effective anti-HCV therapies. Such inhibitors would have therapeutic potential as protease inhibitors, particularly as serine protease inhibitors, and more particularly as HCV NS3 protease inhibitors. Specifically, such compounds may be useful as antiviral agents, particularly as anti-HCV agents.

SUMMARY OF THE INVENTION

The present invention provides compounds, and pharmaceutically acceptable derivatives thereof, that are useful as protease inhibitors, particularly as serine protease inhibitors, and more particularly as HCV NS3 protease inhibitors. These compounds can be used alone or in combination with immunomodulatory agents, such as α -, β - or γ -interferons; other antiviral agents such as ribavirin and amantadine; other inhibitors of hepatitis C protease; inhibitors of other targets in the HCV life cycle including the helicase, polymerase, metalloprotease, or internal ribosome entry; or combinations thereof.

The present invention also provides methods for inhibiting proteases, particularly serine proteases, and more particularly HCV NS3 protease.

The present invention also provides pharmaceutical compositions comprising the compounds of

this invention, as well as multi-component compositions comprising additional immunomodulatory agents, such as α -, β - or γ -interferons; other antiviral agents such as ribavirin and amantadine; other inhibitors of hepatitis C protease; inhibitors of other targets in the HCV life cycle including the helicase, polymerase, metalloprotease, or internal ribosome entry; or combinations thereof. The invention also provides methods of using the compounds of this invention, as well as other related compounds, for the inhibition of HCV.

DETAILED DESCRIPTION OF THE INVENTION

In order that the invention herein described may be more fully understood, the following detailed description is set forth. In the description, the following abbreviations are used:

	<u>Designation</u>	<u>Reagent or Fragment</u>
	Abu	aminobutyric acid
	Ac	acetyl
	AcOH	acetic acid
20	Bn	benzyl
	Boc	tert-butyloxycarbonyl
	Bz	benzoyl
	Cbz	carbobenzyloxy
	CDI	carbonyldiimidazole
25	DCE	1,2-dichloroethane
	DCM	dichloromethane
	DIEA	diisopropylethylamine
	DMA	dimethylacetamide
	DMAP	dimethylaminopyridine
30	DMF	dimethylformamide
	DPPA	diphenylphosphorylazide

	DMSO	dimethylsulfoxide
	Et	ethyl
	EtOAc	ethyl acetate
	Fmoc	9-fluorenylmethoxycarbonyl
5	HbtU	O-benzotriazolyl-N,N,N',N'- tetramethyluronium hexafluorophosphate
	HOBt	N-hydroxybenzotriazole
	HPLC	high performance liquid 10 chromatography
	Me	methyl
	MS	mass spectrometry
	NMP	N-methyl pyrrolidinone
	ND	not determined
15	Pip	piperidine
	Prz	piperazine
	PyBrop	bromo-tris-pyrrolidinophosphonium hexafluorophosphate
	Pyr	pyridine
20	THF	tetrahydrofuran
	TFA	trifluoroacetic acid
	TFE	trifluoroethanol
	Tol	toluene

The following terms are used herein:

25 Unless expressly stated to the contrary, the terms "-SO₂-" and "-S(O)₂-" as used herein refer to a sulfone or sulfone derivative (i.e., both appended groups linked to the S), and not a sulfinic ester.

The term "substituted" refers to the
30 replacement of one or more hydrogen radicals in a given structure with a radical selected from a specified group. When more than one hydrogen radical may be replaced with

a substituent selected from the same specified group, the substituents may be either the same or different at every position.

As used herein, the term "amino" refers to a
5 trivalent nitrogen which may be primary or which may be substituted with 1-2 alkyl groups.

The term "alkyl" or "alkane", alone or in combination with any other term, refers to a straight-chain or branched-chain saturated aliphatic hydrocarbon
10 radical containing the specified number of carbon atoms, or where no number is specified, preferably from 1-10 and more preferably from 1-5 carbon atoms. Examples of alkyl radicals include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-
15 butyl, pentyl, isoamyl, n-hexyl and the like.

The term "alkenyl" or "alkene", alone or in combination with any other term, refers to a straight-chain or branched-chain mono- or poly-unsaturated aliphatic hydrocarbon radical containing the specified
20 number of carbon atoms, or where no number is specified, preferably from 2-10 carbon atoms and more preferably, from 2-5 carbon atoms. Examples of alkenyl radicals include, but are not limited to, ethenyl, E- and Z-propenyl, E- and Z-isobutenyl, E- and Z-pentenyl, E- and
25 Z-hexenyl, E,E-, E,Z-, Z,E-, and Z-Z-hexadienyl and the like.

The term "alkynyl" or "alkyne", alone or in combination with any other term, refers to a straight-chain or branched-chain mono or poly-unsaturated
30 aliphatic hydrocarbon radical containing the specified number of carbon atoms, or where no number is specified, preferably from 2-10 carbon atoms and more preferably,

from 2-5 carbon atoms, wherein at least one of the unsaturated aliphatic hydrocarbon radicals comprises a triple bond. Examples of alkynyl radicals include, but are not limited to, ethynyl, propynyl, isobutynyl, 5 pentynyl, hexynyl, hexenynyl, and the like.

The term "aryl", alone or in combination with any other term, refers to a carbocyclic aromatic radical containing the specified number of carbon atoms, and which may be optionally fused, for example benzofused, 10 with one to three cycloalkyl, aromatic, heterocyclic or heteroaromatic rings. Preferred aryl groups have from 6-14 carbon atoms, and more preferred groups from 6-10 carbon atoms. Examples of aryl radicals include, but are not limited to, phenyl, naphthyl, anthracenyl and the 15 like.

The term "carbocycle", alone or in combination with any other term, refers to a stable non-aromatic 3- to 8-membered carbon ring radical which may be saturated, mono-unsaturated or poly-unsaturated, and which may be 20 optionally fused, for example benzofused, with one to three cycloalkyl, aromatic, heterocyclic or heteroaromatic rings. The carbocycle may be attached at any endocyclic carbon atom which results in a stable structure.

25 The terms "cycloalkyl" or "cycloalkane", alone or in combination with any other term, refers to a stable non-aromatic 3- to 8-membered carbon ring radical which is saturated and which may be optionally fused, for example benzofused, with one to three cycloalkyl, 30 aromatic, heterocyclic or heteroaromatic rings. The cycloalkyl may be attached at any endocyclic carbon atom which results in a stable structure. Preferred

carbocycles have 5 to 6 carbons. Examples of carbocycle radicals include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclopentenyl, cyclohexenyl, indane, 5 tetrahydronaphthalene and the like.

The term "cycloalkenyl" or "cycloalkene" alone or in combination with any other term, refers to a stable cyclic hydrocarbon ring radical containing at least one endocyclic carbon-carbon double bond. The carbocycle may be attached at any cyclic carbon atom which results in a stable structure. Where no number of carbon atoms is specified, a cycloalkenyl radical preferably has from 5-7 carbon atoms. Examples of cycloalkenyl radicals include, but are not limited to, cyclopentenyl, cyclohexenyl, 10 cyclopentadienyl, indenyl and the like.

The term "cycloalkylidenyl", alone or in combination with any other term, refers to a stable cyclic hydrocarbon ring radical containing at least one exocyclic carbon-carbon double bond, wherein the cyclic hydrocarbon ring may be optionally fused, for example benzofused, with one to three cycloalkyl, aromatic, heterocyclic or heteroaromatic rings. The carbocycle may be attached at any cyclic carbon atom, which results in a stable structure. Where no number of carbon atoms is specified, a cycloalkylidenyl radical preferably has from 5-7 carbon atoms. Examples of cycloalkylidenyl radicals include, but are not limited to, cyclopentylidenyl, cyclohexylidenyl, cyclopentenylidenyl and the like. 20

The skilled practitioner would realize that certain groups could be classified either as cycloalkanes or as aryl groups. Examples of such groups include indanyl and tetrahydro naphthyl groups. 30

The term "monocycle" or "monocyclic" alone or in combination with any other term, unless otherwise defined herein, refers to a 5-7 membered ring system.

5 The term "bicycle" or "bicyclic" alone or in combination with any other term, unless otherwise defined herein, refers to a 6-11 membered ring system.

The term "tricycle" or "tricyclic" alone or in combination with any other term, unless otherwise defined herein, refers to a 11-15 membered ring system.

10 The terms "heterocyclyl" and "heterocycle", alone or in combination with any other term, unless otherwise defined herein, refers to a stable 5- to 15-membered mono-, bi-, or tricyclic, heterocyclic ring which is either saturated or partially unsaturated, but
15 not aromatic, and which may be optionally fused, for example benzofused, with one to three cycloalkyl, aromatic, heterocyclic or heteroaromatic rings. Each heterocycle consists of one or more carbon atoms and from one to four heteroatoms selected from the group
20 consisting of nitrogen, oxygen and sulfur. As used herein, the terms "nitrogen and sulfur heteroatoms" include any oxidized form of nitrogen and sulfur, and the quaternized form of any basic nitrogen. A heterocycle may be attached at any endocyclic carbon or heteroatom
25 which results in the creation of a stable structure.

Preferred heterocycles defined above include, for example, imidazolidinyl, indazolinolyl, perhydropyridazyl, pyrrolinyl, pyrrolidinyl, piperidinyl, pyrazolinyl, piperazinyl, morpholinyl, thiamorpholinyl,
30 β -carbolinyl, thiazolidinyl, thiamorpholinyl sulfone, oxopiperidinyl, oxopyrrolidinyl, oxoazepinyl, azepinyl, furazanyl, tetrahydropyranyl, tetrahydrofuranyl,

oxathioly, dithioly, tetrahydrothiophenyl, dioxanyl,
dioxolanyl, tetrahydrofurotetrahydrofuranyl,
tetrahydropyranotetrahydrofuranyl,
tetrahydrofurodihydrofuranyl,
5 tetrahydropyranodihydrofuranyl, dihydropyranyl,
dihydrofuranyl, dihydrofurotetrahydrofuranyl,
dihydropyranotetrahydrofuranyl, sulfolanyl and the like.


The terms "heteroaryl" and "heteroaromatic"
alone or in combination with any other term, unless
10 otherwise defined herein, refers to a stable 3- to 7-
membered monocyclic heterocyclic ring which is aromatic,
and which may be optionally fused, for example,
benzofused, with one to three cycloalkyl, aromatic,
heterocyclic or heteroaromatic rings. Each
15 heteroaromatic ring consists of one or more carbon atoms
and from one to four heteroatoms selected from the group
consisting of nitrogen, oxygen and sulfur. As used
herein, the terms "nitrogen and sulfur heteroatoms"
include any oxidized form of nitrogen and sulfur, and the
20 quaternized form of any basic nitrogen. A heteroaromatic
ring may be attached at any endocyclic carbon or
heteroatom which results in the creation of a stable,
aromatic structure.

Preferred heteroaromatics defined above
25 include, for example, benzimidazolyl, imidazolyl,
quinolyl, isoquinolyl, indolyl, indazolyl, pyridazyl,
pyridyl, pyrrolyl, pyrazolyl, pyrazinyl, quinoxolyl,
pyranyl, pyrimidinyl, pyridazinyl, furyl, thienyl,
triazolyl, thiazolyl, tetrazolyl, benzofuranyl, oxazolyl,
30 benzoxazolyl, isoxazolyl, isothiazolyl, thiadiazolyl,
thiophenyl, oxadiazolyl, oxatriazolyl, thiatriazolyl,

dithiazolyl, dioxazolyl, oxathiazolyl, triazinyl and the like.

The term "halo" refers to a radical of fluorine, chlorine, bromine or iodine. Preferred halogen radicals include fluorine and chlorine.

In chemical formulas, parentheses are used herein to indicate 1) the presence of more than one atom or group bonded to the same atom or group; or 2) a branching point in a chain (i.e., the group or atom immediately before the open parenthesis is bonded directly to the group or atom immediately after the closed parenthesis). An example of the first use is "N(R¹)₂" denoting two R¹ groups bound to the nitrogen atom. An example of the second use is "-C(O)R¹" denoting an oxygen atom and a R¹ bound to the carbon atom, as in the following structure:




As used herein, "B" indicates a boron atom.

Those of skill in the art will realize that certain combinations of moiety choices for variables in the generic structures set forth throughout this application will produce chemically unstable or unfeasible compounds. Such compounds are not intended to be part of the present invention.


The present invention provides compounds that are useful as protease inhibitors, particularly as serine protease inhibitors, and more particularly as HCV NS3 protease inhibitors. As such, they act by interfering with the life cycle of the HCV virus and other viruses

that are dependent upon a serine protease for proliferation. Therefore, these compounds are useful as antiviral agents.

According to one embodiment, the present invention provides a compound of the formula (I):



In these compounds W is selected from:



15 Each R¹ is hydroxy, alkoxy, or aryloxy, or each R¹ is an oxygen atom and together with the boron, to which

they are each bound, form a 5-7 membered ring, wherein the ring atoms are carbon, nitrogen, or oxygen.

Each R^2 is independently hydrogen, halo, alkyl, alkenyl, aryl, aralkyl, aralkenyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, heterocyclyl, heterocyclylalkyl, heterocyclylalkenyl, heteroaryl, or heteroaralkyl, or two R^2 groups, which are bound to the same nitrogen atom, form together with that nitrogen atom, a 5-7 membered monocyclic heterocyclic ring system; wherein any R^2 carbon atom is optionally substituted with J.

Each J is independently alkyl, aryl, aralkyl, alkoxy, aryloxy, aralkoxy, cycloalkyl, cycloalkoxy, heterocyclyl, heterocyclyoxy, heterocyclylalkyl, keto, hydroxy, amino, alkylamino, alkanoylamino, aroylamino, aralkanoylamino, carboxy, carboxyalkyl, carboxamidoalkyl, halo, cyano, nitro, formyl, acyl, sulfonyl, or sulfonamido and is optionally substituted with 1 to 3 J^1 groups.

Each J^1 is independently selected from alkyl, aryl, aralkyl, alkoxy, aryloxy, heterocyclyl, heterocyclyoxy, keto, hydroxy, amino, alkanoylamino, aroylamino, carboxy, carboxyalkyl, carboxamidoalkyl, halo, cyano, nitro, formyl, sulfonyl, or sulfonamido.

L is alkyl, alkenyl, or alkynyl, wherein any hydrogen bound to a carbon atoms is optionally substituted with halogen, and wherein any hydrogen or halogen atom bound to any terminal carbon atom is optionally substituted with sulfhydryl or hydroxy.

Each M is independently alkyl, cycloalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, or heteroaralkyl, and is optionally

substituted by 1 to 3 J groups, wherein any alkyl carbon atom may be replaced by a heteroatom.

R^{18} is a bond, $-N(R^{11})-$, or $-C(O)-$.

R^{11} is hydrogen or C_1 - C_3 alkyl.

5 Each R^{19} is independently H or R^{21} -aryl, or 2 adjacent R^{19} may be bound to one another to form a 5-7 membered aromatic ring, wherein any R^{19} is optionally substituted with 1 to 4 independently selected J' groups. When 2 adjacent R^{19} are bound to one another to form a 5-7
10 membered aromatic ring a bicyclic ring system is formed consisting of the aromatic ring shown in formula I and the aromatic ring formed by two adjacent R^{19} groups. When R^{19} is R^{21} -aryl, this optional substitution may occur on one or more carbon atoms of R^{21} and/or on one or more ring
15 atoms of said aryl. When 2 adjacent R^{19} are bound to one another to form a 5-7 membered aromatic ring, the optional substitution may occur on one or more atoms of the resulting aromatic ring.

 Each R^{21} is independently C_1 - C_3 -straight or
20 branched alkyl, C_2 - C_3 -straight or branched alkenyl, $O-(C_1-C_3)$ -straight or branched alkyl, or $O-(C_2-C_3)$ -straight or branched alkenyl.

n is 0 or 1.


 The ring to which R^{18} and R^{19} are attached may be
25 saturated, partially saturated, aromatic or fully unsaturated. Up to 3 carbon atoms that make up the ring to which R^{18} and R^{19} are attached are optionally replaced with a heteroatom which is independently selected from O, S, $S(O)$, $S(O)_2$ or $N(R^{11})$.

30 A^2 is a bond or $-N(R^{11})-R^{17}(M)-R^{22}-$, wherein R^{17} is $-CH-$ or $-N-$; and R^{22} is $-C(O)-$ or $-S(O)_2-$.

V is a bond, $-\text{CH}(\text{R}^{11})-$, $-\text{O}-$, $-\text{S}-$, or $-\text{N}(\text{R}^{11})-$.

K is a bond, $-\text{O}-$, $-\text{S}-$, $-\text{C}(\text{O})-$, $-\text{S}(\text{O})-$, $-\text{S}(\text{O})_2-$,
or $-\text{S}(\text{O})\text{NR}^{11}-$.

T is $-\text{R}^{12}$, $-\text{alkyl}-\text{R}^{12}$, $-\text{alkenyl}-\text{R}^{12}$, $-\text{alkynyl}-\text{R}^{12}$,
5 $-\text{OR}^{12}$, $-\text{N}(\text{R}^{12})_2$, $-\text{C}(\text{O})\text{R}^{12}$, $-\text{C}(=\text{NO-alkyl})\text{R}^{12}$, or




Each R^{12} is independently selected from
hydrogen, aryl, heteroaryl, cycloalkyl, heterocyclyl,
cycloalkylidenyl, or heterocycloalkylidenyl, and is
10 optionally substituted with 1 to 3 J groups; or a first
 R^{12} and a second R^{12} , together with the nitrogen to which
they are bound, form a mono- or bicyclic ring system
optionally substituted by 1 to 3 J groups.

R^{10} is alkyl, cycloalkyl, aryl, aralkyl,
15 heterocyclyl, heterocyclylalkyl, heteroaryl,
heteroaralkyl, carboxyalkyl, or carboxamidoalkyl, and is
optionally substituted with 1 to 3 J groups.


R^{15} is alkyl, cycloalkyl, aryl, aralkyl,
heterocyclyl, heterocyclylalkyl, heteroaryl,
20 heteroaralkyl, carboxyalkyl, or carboxamidoalkyl, and is
optionally substituted with 1 to 3 J groups.

R^{16} is hydrogen, alkyl, aryl, heteroaryl,
cycloalkyl, or heterocyclyl.


Preferably, W is



5 More preferably, W is



Even more preferably, W is



10 Most preferably, W is -C(O)H.

Preferably, J is alkyl, alkoxy, aryloxy, aryl, aralkyl, aralkoxy, halo, heteroaryl, cyano, amino, nitro, heterocyclyl, acyl, carboxy, carboxyalkyl, alkylamino, hydroxy, heterocyclylalkyl, aralkanoylamino, aroylamino, 5 alkanoylamino, formyl or keto.

More preferably, J is t-butyl, methyl, trifluoromethyl, hydroxy, methoxy, ethoxy, trifluoromethoxy, carboxy, phenyl, benzyl, phenoxy, benzyloxy, fluoro, chloro, bromo, isoxazolyl, pyridinyl, 10 piperidinyl, carboxymethyl, carboxyethyl, dialkylamino, morpholinylmethyl, phenylacetylamino, or acylamino.

Preferably, J¹ is alkoxy, alkyl, halo or aryl.

More preferably, J¹ is C₁₋₃ alkoxy, chloro, C₁₋₃ alkyl, or phenyl.

15 Preferably, L is alkyl, alkenyl, allyl, or propargyl.

More preferably, L is trihalomethyl, sulfhydryl or alkyl substituted with trihalomethyl, sulfhydryl, or hydroxy. Most preferably, L is -CH₂CH₃ or -CH₂CF₃.

20 Preferably, R² is H, fluorine, trifluoromethyl, alkyl, aryl, aralkyl, heteroaralkyl, heterocyclyl, or heterocyclylalkyl. More preferred is when R² is H.

Preferably, M is alkyl, heteroaralkyl, aryl, cycloalkylalkyl, aralkyl, or aralkyl, wherein one of the 25 alkyl carbon atoms is replaced by O or S.

More preferably M is isopropyl, propyl, methyl, pyridylmethyl, benzyl, naphthylmethyl, phenyl, imidazolylmethyl, thiophenylmethyl, cyclohexylmethyl, phenethyl, benzylthiomethyl, or benzyloxyethyl. Most 30 preferably, M is isopropyl.

Preferably, one R¹⁹ is R²¹-aryl and the other two R¹⁹ are H or two R¹⁹ are bound together to form an aromatic

ring and the other R^{19} is H. More preferred is when one R^{19} is $-O-(C_1-C_3)\text{-alkyl-aryl}$ or the two R^{19} bound together form a 6-membered aromatic ring. Most preferred is when one R^{19} is $-O\text{-benzyl}$ or the two R^{19} bound together form phenyl.

Preferably, R^{18} is $-N(R^{11})-$. More preferably, R^{18} is $-N(H)-$ or $-N(CH_3)-$.


Preferably, the ring to which R^{18} and R^{19} are attached is aromatic.

It is preferred that A^2 be a bond or $-N(R^{11})-C(M)-C(O)-$. More preferred is when A^2 is a bond or $-N(H)-C(M)-C(O)-$, wherein M is isopropyl.

Preferably, V is $-N(R^{11})-$. More preferably, V is $-NH-$.

Preferably, K is $-C(O)-$ or $-S(O)_2-$. More preferably, K is $-C(O)-$.

Preferably, T is $-R^{12}$, $-\text{alkyl-}R^{12}$, $-\text{alkenyl-}R^{12}$, $-\text{OR}^{12}$, $-N(R^{12})_2$, $-C(=\text{NO-alkyl})-R^{12}$, or



More preferably, T is $-R^{12}$ or $-\text{alkyl-}R^{12}$.


Preferably, R^{12} is aryl or heteroaryl and is optionally substituted by 1-3 J groups. More preferably, R^{12} is naphthyl, pyrazinyl, or pyridyl, any of which is optionally substituted with a hydroxy group

Preferably, R^{10} is alkyl substituted with carboxy. More preferably, R^{10} is C_{1-3} alkyl substituted with carboxy.








Preferably, R^{15} is alkyl substituted with carboxy. More preferably, R^{15} is C_{1-3} alkyl substituted with carboxy.

The most preferred compounds of this invention are listed in Table 1, below.

Table 1. Preferred Compounds of the formula:



Compound	T	A ²	R ¹⁸	R ^{19A}	R ^{19B}	L
101			-N(H)-	H	O-benzyl	-CH ₂ CH ₃
102			-N(H)-	H	O-benzyl	-CH ₂ CF ₃
103			-N(CH ₃)-	H	O-benzyl	-CH ₂ CH ₃
104		Bond	-N(H)-	H	O-benzyl	-CH ₂ CH ₃
105		Bond	-N(H)-	H	O-benzyl	-CH ₂ CH ₃

Compound	T	A ²	R ^{III}	R ^{IVa}	R ^{IVb}	L
106		Bond	-N(H)-	H	O-benzyl	-CH ₂ CH ₃
107			-N(H)-	H	O-3,4-dichlorobenzyl	-CH ₂ CH ₃
108			-N(H)-	H	H	-CH ₂ CH ₃
109			-N(H)-	bound to one another to form a benzofused ring		-CH ₂ CH ₃

This invention anticipates that many active-site directed inhibitors of the NS3 protease may be peptidomimetic in nature and thus may be designed from the natural substrate. Therefore, preferred substituents in peptidomimetic inhibitors of this invention include those which correspond to the backbone or side chains of naturally occurring substrates or synthetic substrates with high affinity for the enzyme (low K_m).

The skilled practitioner would realize that some certain groups could be classified either as heterocycles or heteroaromatics, depending on the point of attachment.

The compounds of this invention may contain one or more asymmetric carbon atoms and thus may occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers.

All such isomeric forms of these compounds are expressly included in the present invention. Each stereogenic carbon may be of the R or S configuration. Combinations of substituents and variables envisioned by this invention are only those that result in the formation of stable compounds. The term "stable", as used herein, refers to compounds which possess stability sufficient to allow manufacture and which maintains the integrity of the compound for a sufficient period of time to be useful for the purposes detailed herein (e.g., therapeutic or prophylactic administration to a mammal or for use in affinity chromatography applications). Typically, such compounds are stable at a temperature of 40°C or less, in the absence of moisture or other chemically reactive conditions, for at least a week.

The compounds of this invention may be synthesized using conventional techniques. Advantageously, these compounds are conveniently synthesized from readily available starting materials.

As used herein, the compounds of this invention are defined to include pharmaceutically acceptable derivatives or prodrugs thereof. A "pharmaceutically acceptable derivative or prodrug" means any pharmaceutically acceptable salt, ester, salt of an ester, or other derivative of a compound of this invention which, upon administration to a recipient, is capable of providing (directly or indirectly) a compound of this invention.

Accordingly, this invention also provides prodrugs of the compounds of this invention, which are derivatives that are designed to enhance biological properties such as oral absorption, clearance, metabolism

or compartmental distribution. Such derivations are well known in the art.

As the skilled practitioner realizes, the compounds of this invention may be modified by appending
5 appropriate functionalities to enhance selective biological properties. Such modifications are known in the art and include those which increase biological penetration into a given biological compartment (e.g., blood, lymphatic system, central nervous system),
10 increase oral availability, increase solubility to allow administration by injection, alter metabolism and alter rate of excretion.

The term "protected" refers to when the designated functional group is attached to a suitable
15 chemical group (protecting group). Examples of suitable amino protecting groups and protecting groups are described in T.W. Greene and P.G.M. Wuts, Protective Groups in Organic Synthesis, 2d. Ed., John Wiley and Sons (1991); L. Fieser and M. Fieser, Fieser and Fieser's
20 Reagents for Organic Synthesis, John Wiley and Sons (1994); L. Paquette, ed. Encyclopedia of Reagents for Organic Synthesis, John Wiley and Sons (1995) and are exemplified in certain of the specific compounds used in this invention.

25 Particularly favored derivatives and prodrugs are those that increase the bioavailability of the compounds of this invention when such compounds are administered to a mammal (e.g., by allowing an orally administered compound to be more readily absorbed into
30 the blood), have more favorable clearance rates or metabolic profiles, or which enhance delivery of the parent compound to a biological compartment (e.g., the

brain or lymphatic system) relative to the parent species. Preferred prodrugs include derivatives where a group which enhances aqueous solubility or active transport through the gut membrane is appended to the
5 structure of formula (I).

Pharmaceutically acceptable salts of the compounds of this invention include those derived from pharmaceutically acceptable inorganic and organic acids and bases. Examples of suitable acid salts include ace-
10 tate, adipate, alginate, aspartate, benzoate, benzene-sulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptanoate, glycerophosphate, glycolate,
15 hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, palmoate, pectinate, persulfate, 3-phenylpropionate, phosphate,
20 picrate, pivalate, propionate, salicylate, succinate, sulfate, tartrate, thiocyanate, tosylate and undecanoate. Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining
25 the compounds of the invention and their pharmaceutically acceptable acid addition salts.

Salts derived from appropriate bases include alkali metal (e.g., sodium and potassium), alkaline earth metal (e.g., calcium and magnesium), salts with organic
30 bases, such as dicyclohexylamine salts, N-methyl-D-glucamine, salts with amino acids such as arginine and lysine, ammonium and N-(C₁₋₄ alkyl)₄⁺ salts.

This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates, such as dimethyl, diethyl, dibutyl and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides, such as benzyl and phenethyl bromides and others. Water or oil-soluble or dispersible products may be obtained by such quaternization.

In general, compounds of formula (I) are obtained via methods illustrated in the Examples. As can be appreciated by the skilled artisan however the synthetic schemes set forth herein are not intended to comprise a comprehensive list of all means by which the compounds described and claimed in this application may be synthesized. Further methods will be evident to those of ordinary skill in the art. Additionally, the various synthetic steps described above may be performed in an alternate sequence or order to give the desired compounds.

Without being bound by theory, we believe that the compounds of this invention interact either covalently or noncovalently with the active site of the HCV NS3 protease and other serine proteases, inhibiting the ability of such an enzyme to cleave natural or synthetic substrates. Noncovalent interactions are advantageous in that they impart relatively greater specificity of inhibition and will not inhibit other undesirable targets, e.g. cysteine proteases. These compounds will therefore have a greater therapeutic index

when administered to mammals than covalent protease inhibitors, which can interact with a wide range of proteases and cause undesirable toxic effects. In contrast, covalent interactions are advantageous in that
5 they impart greater inhibitory potency allowing lower doses may be administered and thus ameliorating any lack of specificity problems.

The novel compounds of the present invention are excellent inhibitors of proteases, particularly
10 serine proteases, and more particularly HCV NS3 protease inhibitors. Accordingly, these compounds are capable of targeting and inhibiting proteases, particularly serine proteases, and more particularly HCV NS3 proteases. As such, these compounds interfere with the life cycle of
15 viruses, including HCV and are thus useful as antiviral agents. Inhibition can be measured by various methods such as the methods of Example 3.

The term "antiviral agent" refers to a compound or drug which possesses viral inhibitory activity. Such
20 agents include reverse transcriptase inhibitors (including nucleoside and non-nucleoside analogs) and protease inhibitors. Preferably the protease inhibitor is a HCV protease inhibitor.

The term "treating" as used herein refers to
25 the alleviation of symptoms of a particular disorder in a patient or the improvement of an ascertainable measurement associated with a particular disorder. As used herein, the term "patient" refers to a mammal, including a human.

30 Thus, according to another embodiment this invention provides pharmaceutical compositions comprising a compound of formula (I) or a pharmaceutically

acceptable salt thereof; an additional agent selected from, but not including, an immunomodulatory agent, such as α -, β -, or γ -interferon; other antiviral agents, such as ribavarin or amantadine; other inhibitors of HCV protease; inhibitors of other targets in the HCV life cycle such as helicase, polymerase, or metalloprotease; or combinations thereof and any pharmaceutically acceptable carrier, adjuvant or vehicle. An alternate embodiment provides compositions comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof; and a pharmaceutically acceptable carrier, adjuvant or vehicle. Such composition may optionally comprise an additional agent selected from an immunomodulatory agent, such as α -, β -, or γ -interferon; other antiviral agents, such as ribavarin; other inhibitors of HCV protease; inhibitors of the HCV helicase; or combinations thereof.

The term "pharmaceutically acceptable carrier or adjuvant" refers to a carrier or adjuvant that may be administered to a patient, together with a compound of this invention, and which does not destroy the pharmacological activity thereof and is nontoxic when administered in doses sufficient to deliver a therapeutic amount of the compound.

Pharmaceutically acceptable carriers, adjuvants and vehicles that may be used in the pharmaceutical compositions of this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, self-emulsifying drug delivery systems (SEDDS) such as α -tocopherol, polyethyleneglycol 1000 succinate, or TPGS, surfactants used in pharmaceutical dosage forms such as Tweens or other similar polymeric delivery

matrices, serum proteins, such as human serum albumin, gelatin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts
5 or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, polylactic acid, polylactic polyglycollic acid, citric acid, cellulose-
10 based substances, such as HPC and HPMC, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol, wool fat. Cyclodextrins such as α -, β -, and γ -cyclodextrin, or chemically modified derivatives
15 such as hydroxyalkylcyclodextrins, including 2- and 3-hydroxypropyl- β -cyclodextrins, or other solubilized derivatives may also be advantageously used to enhance delivery of compounds of formula (I).

The pharmaceutical compositions of this
20 invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. We prefer oral administration or administration by injection. The pharmaceutical compositions of this invention may contain
25 any conventional non-toxic pharmaceutically-acceptable carriers, adjuvants or vehicles. In some cases, the pH of the formulation may be adjusted with pharmaceutically acceptable acids, bases or buffers to enhance the stability of the formulated compound or its delivery
30 form. The term parenteral as used herein includes subcutaneous, intracutaneous, intravenous, intramuscular, intra-articular, intrasynovial, intrasternal,

intrathecal, intralesional and intracranial injection or infusion techniques.

The pharmaceutical compositions may be in the form of a sterile injectable preparation, for example, as
5 a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to techniques known in the art using suitable dispersing or wetting agents (such as, for example, Tween 80) and suspending agents. The sterile injectable preparation may also be a
10 sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are mannitol, water, Ringer's solution and isotonic sodium chloride
15 solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids, such as oleic acid and its glyceride derivatives are
20 useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant such
25 as those described in Pharmacopeia Helvetica (Ph. Helv.) or a similar alcohol, or carboxymethyl cellulose or similar dispersing agents which are commonly used in the formulation of pharmaceutically acceptable dosage forms such as emulsions and/or suspensions. Other commonly
30 used surfactants such as Tweens or Spans and/or other similar emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of

pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

The pharmaceutical compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, and aqueous suspensions and solutions. In the case of tablets for oral use, carriers which are commonly used include lactose, corn starch, dicalcium phosphate and microcrystalline cellulose (Avicel). Lubricating agents, such as magnesium stearate and talc, are also typically added. For oral administration in a capsule form, useful diluents include lactose, dried corn starch and TPGS, as well as the other diluents used in tablets. For oral administration in a soft gelatin capsule form (filled with either a suspension or a solution of a compound of this invention), useful diluents include PEG400, TPGS, propylene glycol, Labrasol, Gelucire, Transcutol, PVP and potassium acetate. When aqueous suspensions are administered orally, the active ingredient is combined with emulsifying and suspending agents, such as sodium CMC, methyl cellulose, pectin and gelatin. If desired, certain sweetening and/or flavoring and/or coloring agents may be added.

The pharmaceutical compositions of this invention may also be administered in the form of suppositories for rectal administration. These compositions can be prepared by mixing a compound of this invention with a suitable non-irritating excipient which is solid at room temperature but liquid at the rectal temperature and therefore will melt in the rectum to

release the active components. Such materials include, but are not limited to, cocoa butter, beeswax, gelatin, glycerin and polyethylene glycols.

Topical administration of the pharmaceutical compositions of this invention is especially useful when the desired treatment involves areas or organs readily accessible by topical application. For application topically to the skin, the pharmaceutical composition should be formulated with a suitable ointment containing the active components suspended or dissolved in a carrier. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petroleum, white petroleum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax, stearic acid, cetyl stearate, cetyl alcohol, lanolin, magnesium hydroxide, kaolin and water. Alternatively, the pharmaceutical composition can be formulated with a suitable lotion or cream containing the active compound suspended or dissolved in a carrier. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters, wax, cetyl alcohol, 2-octyldodecanol, benzyl alcohol and water. The pharmaceutical compositions of this invention may also be topically applied to the lower intestinal tract by rectal suppository formulation or in a suitable enema formulation. Topically-transdermal patches are also included in this invention.

For ophthalmic use, the pharmaceutical compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as

benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutical compositions may be formulated in an ointment such as petrolatum.

The pharmaceutical compositions of this invention may be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art.

Dosage levels of between about 0.01 and about 100 mg/kg body weight per day, preferably between about 0.5 and about 75 mg/kg body weight per day of the protease inhibitor compounds described herein are useful in a monotherapy for the prevention and treatment of antiviral, particularly anti-HCV mediated disease. Typically, the pharmaceutical compositions of this invention will be administered from about 1 to about 5 times per day or alternatively, as a continuous infusion. Such administration can be used as a chronic or acute therapy. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. A typical preparation will contain from about 5% to about 95% active compound (w/w). Preferably, such preparations contain from about 20% to about 80% active compound.

When the compositions of this invention comprise a combination of a compound of formula (I) and one or more additional therapeutic or prophylactic

agents, both the compound and the additional agent should be present at dosage levels of between about 10 to 100%, and more preferably between about 10 to 80% of the dosage normally administered in a monotherapy regimen.

5 According to one embodiment, the pharmaceutical compositions of this invention comprise an additional immunomodulatory agent. Examples of additional immunomodulatory agents include, but are not limited to, α -, β -, and δ -interferons.

10 According to an alternate embodiment, the pharmaceutical compositions of this invention may additionally comprise an anti-viral agent. Examples of anti-viral agents include, ribavirin and amantadine.

 According to another alternate embodiment, the
15 pharmaceutical compositions of this invention may additionally comprise other inhibitors of HCV protease.

 According to yet another alternate embodiment, the pharmaceutical compositions of this invention may additionally comprise an inhibitor of other targets in
20 the HCV life cycle, such as helicase, polymerase, or metalloprotease.

 Upon improvement of a patient's condition, a maintenance dose of a compound, composition or combination of this invention may be administered, if
25 necessary. Subsequently, the dosage or frequency of administration, or both, may be reduced, as a function of the symptoms, to a level at which the improved condition is retained when the symptoms have been alleviated to the desired level, treatment should cease. Patients may,
30 however, require intermittent treatment on a long-term basis upon any recurrence of disease symptoms.

As the skilled artisan will appreciate, lower or higher doses than those recited above may be required. Specific dosage and treatment regimens for any particular patient will depend upon a variety of factors, including
5 the activity of the specific compound employed, the age, body weight, general health status, sex, diet, time of administration, rate of excretion, drug combination, the severity and course of the infection, the patient's disposition to the infection and the judgment of the
10 treating physician.

When these compounds or their pharmaceutically acceptable salts are formulated together with a pharmaceutically acceptable carrier, the resulting composition may be administered in vivo to mammals, such
15 as man, to inhibit serine proteases, particularly HCV NS3 protease or to treat or prevent viral infection, particularly HCV virus infection. Such treatment may also be achieved using the compounds of this invention in combination with agents which include, but are not
20 limited to: immunomodulatory agents, such as α -, β -, or γ -interferons; other antiviral agents such as ribavirin, amantadine; other inhibitors of HCV NS3 protease; inhibitors of other targets in the HCV life cycle such as helicase, polymerase, metalloprotease, or internal
25 ribosome entry; or combinations thereof. These additional agents may be combined with the compounds of this invention to create a single dosage form. Alternatively these additional agents may be separately administered to a mammal as part of a multiple dosage
30 form.

Accordingly, another embodiment of this invention provides methods of inhibiting serine protease

activity in mammals by administering a compound of the formula (I), wherein the substituents are as defined above. Preferably, the serine protease is HCV NS3.

In an alternate embodiment, the invention
5 provides methods of inhibiting HCV or HCV NS3 activity in a mammal comprising the step of administering to said mammal, a compound of formula (I), wherein the substituents are as defined above.

In an alternate embodiment, this invention
10 provides methods of decreasing serine protease activity in a mammal comprising the step of administering to said mammal any of the pharmaceutical compositions and combinations described above. If the pharmaceutical composition comprises only a compound of this invention
15 as the active component, such methods may additionally comprise the step of administering to said mammal an agent selected from an immunomodulatory agent, an antiviral agent, a HCV protease inhibitor, or an inhibitor of other targets in the HCV life cycle. Such
20 additional agent may be administered to the mammal prior to, concurrently with, or following the administration of the HCV inhibitor composition.

In a preferred embodiment, these methods are useful in decreasing HCV NS3 protease activity in a
25 mammal. If the pharmaceutical composition comprises only a compound of this invention as the active component, such methods may additionally comprise the step of administering to said mammal an agent selected from an immunomodulatory agent, an antiviral agent, a HCV
30 protease inhibitor, or an inhibitor of other targets in the HCV life cycle such as helicase, polymerase, or metalloprotease. Such additional agent may be

administered to the mammal prior to, concurrently with, or following the administration of the compositions of this invention.

In an alternate preferred embodiment, these
5 methods are useful for inhibiting viral replication in a mammal. Such methods are useful in treating or preventing, for example, viral diseases, such as HCV. If the pharmaceutical composition comprises only a compound of this invention as the active component, such methods
10 may additionally comprise the step of administering to said mammal an agent selected from an immunomodulatory agent, an antiviral agent, a HCV protease inhibitor, or an inhibitor of other targets in the HCV life cycle. Such additional agent may be administered to the mammal
15 prior to, concurrently with, or following the administration of the composition according to this invention.

The compounds set forth herein may also be used as laboratory reagents. The compounds of this invention
20 may also be used to treat or prevent viral contamination of materials and therefore reduce the risk of viral infection of laboratory or medical personnel or patients who come in contact with such materials. These materials include, but are not limited to, biological materials,
25 such as blood, tissue, etc; surgical instruments and garments; laboratory instruments and garments; and blood collection apparatuses and materials.

In order that this invention be more fully understood, the following examples are set forth. These
30 examples are for the purpose of illustration only and are not to be construed as limiting the scope of the invention in any way.

General Materials and Methods

Compounds 101 through 109 were prepared using the generic synthesis scheme 1, depicted below. Synthesis of specific compounds are described in the following examples.

The correct $(M + H)^+$ and/or $(M + Na)^+$ molecular ions for all compounds as set forth in Table 2 were obtained by either matrix-assisted laser desorption mass spectrometry (Kratos MALDI I) or by electro spray mass spectrometry (MICROMASS Quatro II).

Numerous amino acids for use in the synthesis of peptidyl and peptidomimetic compounds of this invention may be purchased commercially from, for instance, Sigma Chemical Company or Bachem Feinchemikalien AG (Switzerland). Amino acids that are not commercially available can be made by known synthetic routes ("Kinetic Resolution of Unnatural and Rarely Occurring Amino Acids: Enantioselective Hydrolysis of N-Acyl Amino Acids Catalyzed by Acylase I", Chenault, H.K. et. al., J. Am. Chem. Soc. 111, 6354-6364 (1989) and references cited therein; "Synthesis of β - γ -Unsaturated Amino Acids by the Strecker Reaction, Greenlee, W.J., J. Org. Chem. 49, 2632-2634 (1984); "Recent Stereoselective Synthetic Approaches to Beta-amino Acids", Cole, D. Tetrahedron 50: 9517 (1994); "The Chemistry of Cyclic Alpha Imino Acids", Mauger, A.B; Volume 4 of "Chemistry and Biochemistry of Amino Acids, Peptides, and Proteins", Weinstein, B. editor, Marcel Dekker (1977); "Recent Progress in the Synthesis and Reactions of Substituted Piperidines", Org. Prep. Procedure Int. 24, 585-621 (1992), all of which are incorporated herein by reference).






Certain compounds of formula (I) may be synthesized from amino acids by procedures which are well known in the art of peptide and organic chemical synthesis. Examples of such syntheses are generally set forth in Bodanszky and Bodanszky, "The Practice of Peptide Synthesis", Springer-Verlag, Berlin, Germany (1984), "The Peptides", Gross and Meinhofer, eds.; Academic Press, 1979, Vols. I-III, and Stewart, J.M. and Young, J.D., "Solid Phase Peptide Synthesis, Second Edition", Pierce Chemical Company, Rockford, IL (1984); and "Recent Advances in the Generation of Molecular Diversity", Moos, W.H., Green, G.D. and Pavia, M.R. in "Annual Reports in Medicinal Chemistry, Vol. 28" pp. 315-324; Bristol, J.A., ed.; Academic Press, San Diego, CA (1993), all of which are incorporated herein by reference.


Typically, for solution phase synthesis of peptides, the α -amine of the amino acid to be coupled is protected by a urethane such as Boc, Cbz, Fmoc or Alloc while the free carboxyl is activated by reaction with a carbodiimide such as DCC, EDC, or DIC, optionally in the presence of a catalyst such as HOBT, HOAt, HOSu, or DMAP. Other methods, which proceed through the intermediacy of activated esters, acid halides, enzyme-activated amino acids and anhydrides including phosphonium reagents such as BOP, Py-BOP, N-carboxy-anhydrides, symmetrical anhydrides, mixed carbonic anhydrides, carbonic-phosphinic and carbonic-phosphoric anhydrides, are also suitable. After the peptide has been formed, protecting groups may be removed by methods described in the references listed above, such as by hydrogenation in the presence of a palladium, platinum or rhodium catalyst,

treatment with sodium in liquid ammonia, hydrochloric, hydrofluoric, hydrobromic, formic, trifluoromethanesulfonic, or trifluoroacetic acid, secondary amines, fluoride ion, trimethylsilyl halides including bromide and iodide, or alkali. Automation of the synthetic process, using techniques such as those set forth above, can be accomplished by use of commercially available instrumentation, including but not limited to the Advanced Chemtech 357 FBS and 496 MOS; Tecan CombiTec, and Applied Biosystems 433A among others. Specific application of these methods and their equivalents, depending upon the target compound, will be apparent to those skilled in the art. Modifications of chemical processes and choice of instrumentation is within the skill of the ordinary practitioner.

Although some of the schemes depicted below indicate particular stereochemistry for certain groups, it should be apparent to those of skill in the art that the synthesis schemes may be modified to allow for the use of those certain groups having the opposite stereochemistry. Therefore, the indication of stereochemistry in these scheme is not intended to limit the depicted synthesis to any particular stereochemistry of any intermediate or final product.

25


SCHEME 1. General Syntheses for the Anthranilic Acids



EXAMPLE 1Synthesis of Compound 101

Synthesis of 2. To a solution of Boc-5-hydroxyanthranilic acid (1) (1.5 g, 5.8 mmol) in 45 mL of anhydrous THF at room temperature was added portionwise, sodium hydride (581 mg, 14.5 mmol). The mixture was stirred and heated to reflux for 2 hours and cooled to room temperature. Benzyl bromide (691 mL, 5.8 mmol) was then added and the reaction mixture was heated to reflux for 1.5 hours, cooled to room temperature and stirred overnight. The reaction was quenched with ice-water (40 mL) and carefully acidified to pH 3 with HCl 2N. Extraction with ethyl acetate (100 mL) followed by a water (50 mL) and a brine (40 mL) wash gave, after drying over Na₂SO₄ and concentration in *vacuo*, a solid residue. Flash chromatography of the crude mixture (10% methanol-90% dichloromethane) provided 1.42 g (70%) of the desired product 2.

Synthesis of 4. 4-Methyl Benzhydrylamine resin (1.05 mmol/g, 20.0 g) was placed in a sintered glass funnel and washed with dimethylformamide (3 X 75 mL), 10% (v/v) diisopropylethylamine (DIEA) in dimethylformamide (2 X 75 mL) and finally with dimethylformamide (4 X 75 mL). Sufficient dimethylformamide was added to the resin to obtain a slurry followed by:



(8.0 g, 20.8 mmol, prepared from (2S) 2-(t-Butyloxycarbonylamino)-butyraldehyde according to A.M.

Murphy et. al. J. Am. Chem. Soc., 114, 3156-3157 (1992)),
1-hydroxybenzotriazole hydrate (HOBT.H₂O; 3.22 g, 21.0
mmol), O-benzotriazole-N,N,N',N'-tetramethyluronium
hexafluorophosphate (HBTU; 8.0 g, 21.0 mmol), and DIEA
5 (11.0 mL, 63 mmol). The reaction mixture was agitated
overnight at room temperature using a wrist arm shaker.
The resin was isolated on a sintered glass funnel by
suction filtration and washed with dimethylformamide (3 X
75 mL). Unreacted amine groups were then capped by
10 reacting the resin with 20% (v/v) acetic
anhydride/dimethylformamide (2 X 50 mL) directly in the
funnel (10 min/wash). The resin was washed with
dimethylformamide (3 X 75 mL) and dichloromethane (3 X 75
mL) prior to drying overnight *in vacuo* to yield an
15 intermediate resin (26.3 g, 81% yield).

The t-Boc protecting group was removed from the
intermediate resin using the Advanced ChemTech 396
Multiple Peptide synthesizer by the following procedure.
The intermediate resin (0.05 mmol) was swelled by washing
20 with dichloromethane (3 X 1 mL) followed by cleavage of
the t-Boc protecting group with 50% (v/v)
TFA/dichloromethane (1.0 mL) for 10 minutes (with
shaking) followed by fresh reagent (1 mL) for 30 minutes.
The resin was then washed with dichloromethane (3 X 1
25 mL), followed by DMF (3 X 1 mL), then 10%
DIEA/dimethylformamide (v/v) (2 X 1 mL), and finally with
N-methylpyrrolidone (3 X 1 mL) to yield resin 4.

Synthesis of 5. Resin 4 (554 mg, 0.36 mmol) was
placed in a Nalgene syringe equipped with a filter at the
30 bottom and coupled to 2 (251 mg, 0.72 mmol) with HOBT
(110 mg, 0.72 mmol), HBTU (273 mg, 0.72 mmol) and DIEA
(376 mL, 2.16 mmol) in 4 mL of NMP (with shaking) for 72

hours. The solvent was removed by suction and the resin washed sequentially with NMP (3 X 4 mL) and dichloromethane (3 X 4 mL) to provide resin 5.

Synthesis of 6. The t-Boc protecting group was removed from the resin 5 (in the Nalgene syringe) with 50% (v/v) TFA/dichloromethane (6 mL) for 20 min. (with shaking). The resin was washed with dichloromethane (3 X 5 mL), followed by 10% DIEA/dichloromethane (3 X 5 mL) and finally with dichloromethane (3 X 5 mL). To the resulting resin was added Fmoc-valine-N-carboxy anhydride (526 mg, 1.44 mmol) and 6 mL of NMP. The reaction mixture was agitated for 72 hours at room temperature and the solvent was removed by suction. The resin 6 was washed sequentially with NMP (3 X 5 mL) and dichloromethane (3 X 5 mL) and used directly for the next step.

Synthesis of 8. This compound was prepared from resin 6 (0.1 mmol) using an Applied Biosystems Model 433A Peptide synthesizer. N^o-Fmoc-protected amino acid and pyrazine-2-carboxylic acid were added sequentially to resin 6 with standard coupling cycles using HBTU with HOBT as coupling agents in NMP to yield resin 8.

Syntheses of Compound 101. The aldehyde was cleaved from the resin 8 by treatment with 10 mL of a solution of THF/30% formalin/AcOH/0.1N HCl 5:1:1:1 (v:v:v:v) for 1 hour at room temperature. After washing the resin with cleavage reagent (5 mL), the combined filtrates were diluted with water (15 mL) and 5 mL of that mixture was purified by RP-HPLC with a Waters DeltaPak 300 A C18 column (15 m, 30 X 300 mm) eluting with a linear

acetonitrile gradient containing 0.1% TFA (v/v) over 45 min. at 20 mL/min. Fractions containing the desired product were pooled and lyophilized to provide 101.
M + H=617.3.

5

EXAMPLE 2Synthesis of Compound 102

Intermediates 1 through 4 were prepared as in Example 1.

Synthesis of 5. Resin 4 (W=CF₃) (1.2g, .384 mmol) was placed in a Nalgene syringe equipped with a filter at the bottom and coupled to 2 (267 mg, 0.768 mmol) with HOBT (118 mg, 0.768 mmol), HBTU (291 mg, 0.768 mmol) and DIEA (401 mL, 2.3 mmol) in 4 mL of NMP (with shaking) for 72 hours. The solvent was removed by suction and the resin washed sequentially with NMP (3 X 4 mL) and dichloromethane (3 X 4 mL) to provide resin 5.

Synthesis of 6. The t-Boc protecting group was removed from the resin 5 (in the Nalgene syringe) with 50% (v/v) TFA/dichloromethane (6 mL) for 20 minutes (with shaking). The resin was washed with dichloromethane (3 X 5 mL), followed by 10% DIEA/dichloromethane (3 X 5 mL) and finally with dichloromethane (3 X 5 mL). To the resulting resin was added Fmoc-valine-N-carboxy anhydride (562 mg, 1.54 mmol) and 8 mL of NMP. The reaction mixture was agitated for 72 hours at room temperature and the solvent was removed by suction. The resin 6 was washed sequentially with NMP (3 X 6 mL) and dichloromethane (3 X 6 mL) and used directly for the next step.

Synthesis of 8. This compound was prepared from resin 6 (0.1 mmol) using an Applied Biosystems Model 433A Peptide synthesizer. Na-Fmoc-protected amino acid and pyrazine-2-carboxylic acid were added sequentially to resin 6 with standard coupling cycles using HBTU with HOBt as coupling agents in NMP to yield resin 8.

Syntheses of 102. The aldehyde was cleaved from the resin 8 by treatment with 10 mL of a solution of THF/30% formalin/AcOH/0.1N HCl 5:1:1:1 (v:v:v:v) for 1 hour at room temperature. After washing the resin with cleavage reagent (5 mL), the combined filtrates were diluted with water (15 mL) and 5 mL of that mixture was purified by RP-HPLC with a Waters DeltaPak 300 A C18 column (15 m, 30 X 300 mm) eluting with a linear acetonitrile gradient containing 0.1% TFA (v/v) over 45 min. at 20 mL/min. Fraction containing the desired product were pooled and lyophilized to provide 102.
M + H = 671.2

EXAMPLE 3

Synthesis of Compound 103

Synthesis of 3 (R₂ = Me). To a solution of 2 (350 mg, 0.001 mol) in 10 mL of anhydrous DMF at room temperature was added, portionwise, NaH (100 mg, .0025 mol). The mixture was stirred for 40 minutes, then methyl iodide (0.175 mL, 0.0028 mol) was added and the reaction mixture was stirred overnight. The reaction was quenched with water and extracted with ethyl acetate. The organic phase was dried (Na₂SO₄) and concentrated in vacuo to provide 335 mg (89%) of an oil that was used directly for the next step.

To a solution of the above oil (335 mg, 0.0089 mol) in 8 mL of methanol room temperature was added 2.2 mL of 2N NaOH. The reaction was stirred overnight at room temperature after which the methanol was removed under vacuum. The resulting aqueous phase was diluted with water then slowly acidified to pH 3 with 1N HCl. The aqueous phase was extracted with ethyl acetate and the organic phase dried (Na₂SO₄) and concentrated in vacuo to give 300 mg (93%) of 3 that was used directly for the next step.

Synthesis of 6. The t-Boc protecting group was removed from the resin 5 (in the Nalgene syringe) with 50% (v/v) TFA/dichloromethane (6 mL) for 20 min. (with shaking). The resin was washed with dichloromethane (3 X 5 mL), followed by 10% DIEA/dichloromethane (3 X 5 mL) and finally with dichloromethane (3 X 5 mL). To the resulting resin was added Fmoc-valine-N-carboxy anhydride (606 mg, 1.66 mmol) and 10 mL of NMP. The reaction mixture was agitated for 72 hours at room temperature and the solvent was removed by suction. The resin 6 was washed sequentially with NMP (3 X 6 mL) and dichloromethane (3 X 6 mL) and used directly for the next step.

Synthesis of 8. This compound was prepared from resin 6 (0.1 mmol) using an Applied Biosystems Model 433A Peptide synthesizer. Na-Fmoc-protected amino acid and pyrazine-2-carboxylic acid were added sequentially to resin 6 with standard coupling cycles using HBTU with HOBT as coupling agents in NMP to yield resin 8.

Syntheses of 103. The aldehyde was cleaved from the resin 8 by treatment with 10 mL of a solution of THF/30% formalin/AcOH/0.1N HCl 5:1:1:1 (v:v:v:v) for 1 hour at room temperature. After washing the resin with cleavage reagent (5 mL), the combined filtrates were diluted with water (15 mL) and 5 mL of that mixture was purified by RP-HPLC with a Waters DeltaPak 300 A C18 column (15 m, 30 X 300 mm) eluting with a linear acetonitrile gradient containing 0.1% TFA (v/v) over 45 min. at 20 mL/min. Fraction containing the desired product were pooled and lyophilized to provide 103.

M + H = 631.3.

EXAMPLE 4

Synthesis of Compound 104

Intermediates 1 through 6 were prepared as in Example 1.

Synthesis of 7. This compound was prepared from resin 6 (0.1 mmol) using an Applied Biosystems Model 433A Peptide synthesizer. 2-Hydroxynaphthoic acid was added to resin 6 with standard coupling cycles using HBTU with HOBt as coupling agents in NMP to yield resin 7.

Syntheses of 104. The aldehyde was cleaved from the resin 7 by treatment with 3 mL of a solution of THF/30% formalin/AcOH/0.1N HCl 5:1:1:1 (v:v:v:v) for 1 hour at room temperature. After washing the resin with cleavage reagent (1.5 mL), the combined filtrates were diluted with water (4.5 mL) and 5 mL of that mixture was purified by RP-HPLC with a Waters DeltaPak 300 A C18 column (15 m, 30 X 300 mm) eluting with a linear acetonitrile gradient containing 0.1% TFA (v/v) over 45 min. at 20 mL/min.

Fraction containing the desired product were pooled and lyophilized to provide 104.

M + H = 582.2.

EXAMPLE 5

5

Synthesis of Compound 105

Intermediates 1 through 6 were prepared as in Example 1.

Synthesis of 7. This compound was prepared from resin 6 (0.1 mmol) using an Applied Biosystems Model 433A
10 Peptide synthesizer. 2-Hydroxy-4-nicotinic acid was added to resin 6 with standard coupling cycles using HBTU with HOBT as coupling agents in NMP to yield resin 7.

Syntheses of 105. The aldehyde was cleaved from the resin 8 by treatment with 3 mL of a solution of THF/30%
15 formalin/AcOH/0.1N HCl 5:1:1:1 (v:v:v:v) for 1 hour at room temperature. After washing the resin with cleavage reagent (1.5 mL), the combined filtrates were diluted with water (4.5 mL) and 5 mL of that mixture was purified by RP-HPLC with a Waters DeltaPak 300 A C18 column (15 m,
20 30 X 300 mm) eluting with a linear acetonitrile gradient containing 0.1% TFA (v/v) over 45 min. at 20 mL/min.

Fraction containing the desired product were pooled and lyophilized to provide 105.

M + H = 533.2

EXAMPLE 6Synthesis of Compound 106

Intermediates 1 through 6 were prepared as in Example 1.

5 **Synthesis of 7.** This compound was prepared from resin 6 (0.1 mmol) using an Applied Biosystems Model 433A Peptide synthesizer. 6-Hydroxynaphthoic acid was added to resin 6 with standard coupling cycles using HBTU with HOBT as coupling agents in NMP to yield resin 7.

10 **Syntheses of 106.** The aldehyde was cleaved from the resin 8 by treatment with 3 mL of a solution of THF/30% formalin/AcOH/0.1N HCl 5:1:1:1 (v:v:v:v) for 1 hour at room temperature. After washing the resin with cleavage reagent (1.5 mL), the combined filtrates were diluted
15 with water (4.5 mL) and 5 mL of that mixture was purified by RP-HPLC with a Waters DeltaPak 300 A C18 column (15 m, 30 X 300 mm) eluting with a linear acetonitrile gradient containing 0.1% TFA (v/v) over 45 min. at 20 mL/min. Fraction containing the desired product were pooled and
20 lyophilized to provide 106.
M + H = 582.2.

EXAMPLE 7Synthesis of Compound 107

Synthesis of 2. To a solution of Boc-5-
25 hydroxyanthranilic acid (1) (1.5 g, 5.8 mmol) in 45 mL of anhydrous THF at room temperature was added portionwise, sodium hydride (581 mg, 14.5 mmol). The mixture was stirred and heated to reflux for 2 hours and cooled to room temperature. 3,4-dichloro benzyl bromide

(1.39 g, 5.8 mmol) was then added and the reaction mixture was heated to reflux for 1.5 hours, cooled to room temperature and stirred overnight. The reaction was quenched with ice-water (40 mL) and carefully acidified to pH 3 with 2N HCl. Extraction with ethyl acetate (100 mL) followed by a water (50 mL) and a brine (40 mL) wash gave, after drying (Na₂SO₄) and concentration *in vacuo*, a solid residue. Subjection of the crude mixture to flash chromatography (10% methanol-90% dichloromethane) provided 1.37 g (57%) of the desired product 2.

Synthesis of 5. Resin 4 (1.0 g, 0.65 mmol) (from Example 1) was placed in a Nalgene syringe equipped with a filter at the bottom and coupled to 2 (542 mg, 1.3 mmol) with HOBt (199 mg, 1.3 mmol), HBTU (493 mg, 1.3 mmol) and DIEA (679 mL, 3.9 mmol) in 8 mL of NMP (with shaking) for 72 hours. The solvent was removed by suction and the resin washed sequentially with NMP (3 X 8 mL) and dichloromethane (3 X 8 mL) to provide resin 5.

Synthesis of 6. The *t*-Boc protecting group was removed from the resin 5 (in the Nalgene syringe) with 25% (v/v) TFA/dichloromethane (10 mL) for 30 minutes (with shaking). The resin was washed with dichloromethane (3 X 7 mL), followed by 10% DIEA/dichloromethane (3 X 7 mL) and finally with dichloromethane (3 X 7 mL). To the resulting resin was added Fmoc-valine-N-carboxy anhydride (950 mg, 2.6 mmol) and 8 mL of NMP. The reaction mixture was agitated for 72 hours at room temperature and the solvent was removed by suction. The resin 6 was washed sequentially with NMP (3 X 7 mL) and dichloromethane (3 X 7 mL) and used directly for the next step.

Synthesis of 8. This compound was prepared from resin 6 (0.1 mmol) using an Applied Biosystems Model 433A Peptide synthesizer. N^α-Fmoc-protected amino acid and pyrazine-2-carboxylic acid were added sequentially to resin 6 with standard coupling cycles using HBTU with HOBt as coupling agents in NMP to yield resin 8.

Syntheses of 107. The aldehyde was cleaved from the resin 8 by treatment with 10 mL of a solution of THF/30% formalin/AcOH/0.1N HCl 5:1:1:1 (v:v:v:v) for 1 hour at room temperature. After washing the resin with cleavage reagent (5 mL), the combined filtrates were diluted with water (15 mL) and 5 mL of that mixture was purified by RP-HPLC with a Waters DeltaPak 300 A C18 column (15 m, 30 X 300 mm) eluting with a linear acetonitrile gradient containing 0.1% TFA (v/v) over 45 minutes at 20 mL/minute. Fraction containing the desired product were pooled and lyophilized to provide 107.

M + H = 685.1.

EXAMPLE 8

Synthesis of Compound 108

Synthesis of 5. Resin 4 (640 mg, 0.414 mmol) (from Example 1) was placed in a Nalgene syringe equipped with a filter at the bottom and coupled to 1 (wherein X, Y and Z are hydrogen; 200 mg, 0.828 mmol) with HOBt (126 mg, 0.828 mmol), HBTU (314 mg, 0.828 mmol) and DIEA (433 mL, 3.9 mmol) in 6 mL of NMP (with shaking) for 24 h. The solvent was removed by suction and the resin washed sequentially with NMP (3 X 5 mL) and dichloromethane (3 X 5 mL) to provide resin 5.

Synthesis of 6. The t-Boc protecting group was removed from the resin 5 (in the Nalgene syringe) with 50% (v/v) TFA/dichloromethane (6 mL) for 20 minutes (with shaking). The resin was washed with 5 dichloromethane (3 X 6 mL), followed by 10% DIEA/dichloromethane (3 X 6 mL) and finally with dichloromethane (3 X 6 mL). To the resulting resin was added Fmoc-valine-N-carboxy anhydride (605 mg, 1.66 mmol) and 10 mL of NMP. The reaction mixture was agitated for 10 72 hours at room temperature and the solvent was removed by suction. The resin 6 was washed sequentially with NMP (3 X 7 mL) and dichloromethane (3 X 7 mL) and used directly for the next step.

Synthesis of 8. This compound was prepared from 15 resin 6 (0.1 mmol) using an Applied Biosystems Model 433A Peptide synthesizer. N^α-Fmoc-protected amino acid and pyrazine-2-carboxylic acid were added sequentially to resin 6 with standard coupling cycles using HBTU with HOBT as coupling agents in NMP to yield resin 8.

20 Syntheses of 108. The aldehyde was cleaved from the resin 8 by treatment with 10 mL of a solution of THF/30% formalin/AcOH/0.1N HCl 5:1:1:1 (v:v:v:v) for 1 hour at room temperature. After washing the resin with cleavage reagent (5 mL), the combined filtrates were diluted with 25 water (15 mL) and 5 mL of that mixture was purified by RP-HPLC with a Waters DeltaPak 300 A C18 column (15 m, 30 X 300 mm) eluting with a linear acetonitrile gradient containing 0.1% TFA (v/v) over 45 minutes at 20 mL/minute. Fraction containing the desired product were 30 pooled and lyophilized to provide 108.
M + H = 511.3.

EXAMPLE 9Synthesis of Compound 109

Synthesis of 5. Resin 4 (640 mg, 0.414 mmol) (from Example 1) was placed in a Nalgene syringe equipped with a filter at the bottom and coupled to 1 (238 mg, 0.828 mmol) with HOBt (126 mg, 0.828 mmol), HBTU (314 mg, 0.828 mmol) and DIEA (433 mL, 3.9 mmol) in 6 mL of NMP (with shaking) for 24 hours. The solvent was removed by suction and the resin washed sequentially with NMP (3 X 5 mL) and dichloromethane (3 X 5 mL) to provide resin 5.

Synthesis of 6. The t-Boc protecting group was removed from the resin 5 (in the Nalgene syringe) with 50% (v/v) TFA/dichloromethane (6 mL) for 20 minutes (with shaking). The resin was washed with dichloromethane (3 X 6 mL), followed by 10% DIEA/dichloromethane (3 X 6 mL) and finally with dichloromethane (3 X 6 mL). To the resulting resin was added Fmoc-valine-N-carboxy anhydride (605 mg, 1.66 mmol) and 10 mL of NMP. The reaction mixture was agitated for 72 hours at room temperature and the solvent was removed by suction. The resin 6 was washed sequentially with NMP (3 X 7 mL) and dichloromethane (3 X 7 mL) and used directly for the next step.

Synthesis of 8. This compound was prepared from resin 6 (0.1 mmol) using an Applied Biosystems Model 433A Peptide synthesizer. N^α-Fmoc-protected amino acid and pyrazine-2-carboxylic acid were added sequentially to resin 6 with standard coupling cycles using HBTU with HOBt as coupling agents in NMP to yield resin 8.


$$M + H = 561.3.$$

Synthesis of Para-Hydroxyanthranilic Acid Derivatives

NC(=O)C1=CC=C(C=C1)C(=O)NC(CW)C=O

9a

or



is achieved using the steps set forth in Scheme 1 to produce 9 or 10 and intermediate 1 containing a hydroxy group at Y and hydrogens at X and Z.





Synthesis of 1. This compound is prepared in three steps from 2-nitro-4-aminobenzoic acid. Diazotization of 2-Nitro-4-aminobenzoic with, for example, sodium nitrite in 20% sulfuric acid provides 2-nitro-4-hydroxybenzoic acid. Reduction of 2-nitro-4-hydroxybenzoic acid with, for example, tin in concentrated hydrochloric acid provide the desired *p*-hydroxyanthranilic acid. The amino


group is then protected, for example, by treatment with, for example Boc anhydride in dichloromethane with diisopropylethylamine to give 1.

EXAMPLE 11

5 Scheme 2, set forth below, depicts the synthetic route to obtain alkyl-substituted anthranilic acid derivative compounds of this invention.

SCHEME 2. General Syntheses for the Anthranilic Acids







Synthesis of 12. This compound is prepared in three steps from the coupling of 11 (wherein X is Br; Y and Z are H) with styrene in a Heck reaction using palladium acetate and triethylamine. After deprotection and hydrogenation of the double bond with palladium on carbon followed by protection of the amino group as a carbamate


with Boc anhydride in dichloromethane and DIEA compound 12 is obtained.

Para-alkyl-substituted anthranilic acid derivatives are similarly prepared using 11 wherein Y is
5 Br and X and Z are H, similar to the method described in this example.


EXAMPLE 12

Carbazate-containing compounds of this invention (compounds 27 and 28) are produced according to
10 Scheme 3, depicted below:






25




26

THF / H₂CO / AcOH /
0.1N HCl



27

THF / H₂CO / AcOH /
0.1N HCl



28

Synthesis of 22. This compound could be prepared for example, in two steps from 4-hydroxyphthalic acid (20) by alkylation with, for example, an excess of benzyl bromide with a base such as potassium carbonate in dimethylformamide. After saponification of the diester 21 with, for example, potassium hydroxide in ethanol provide compound 22.

10

Synthesis of 23 and 24. These compounds could be prepared by coupling 22 with pyrazine-Val-isopropylcarbazate or with arylcarboxylic acid (Cap) isopropylcarbazate with, for example, HOBt, HBTU and DIEA


in N-methylpyrrolidinone as previously described in Example 1 to provide 23 or 24.

Synthesis of 25 and 26. These resins could be
5 prepared from resin 4 according to Scheme 3 and
experimental procedures previously described in Example 1.

Syntheses of 27 and 28. These compounds could be
prepared according to Scheme 2 and experimental
10 procedures previously described in Examples 1 and 2.

EXAMPLE 13

1,4-diketone-containing compounds of this
invention (compounds 32 and 33) are produced according to
15 Scheme 4, depicted below:

SCHEME 4. General Syntheses for the 1,4-Diketones


Synthesis of 30. This compound is prepared from 29 by deprotonation/halogen-metal exchange with a base, such as butyl lithium in tetrahydrofuran, followed by addition of Boc-Val-C(O)N(OMe)Me (Weinreb amide).


Syntheses of 32 and 33. These compounds are prepared according to Scheme 4 and experimentals previously described in Examples 1 and 2.

EXAMPLE 14

5 1,2-diketone-containing compounds of this invention (compounds 39 and 40) are produced according to Scheme 5, depicted below:



SCHEME 5. General Syntheses for the 1,2-Diketones







1) 50% TFA-DCM
2) CapCO₂H / HBTU / HOBT / DIEA / NMP

1) 50% TFA-DCM
2) Fmoc-Val / HBTU / HOBT / DIEA / NMP
3) 25% pip / DMF
4) Pyr-OH / HBTU / HOBT / DIEA / NMP



THF / H₂CO / AcOH /
0.1N HCl

THF / H₂CO / AcOH /
0.1N HCl



Synthesis of 35. This compound is prepared from phosphonium bromide 34 by treatment with a base such as potassium hexamethyldisilazide in tetrahydrofuran, followed by addition of Boc-Val-H.


Synthesis of 36. This compound is prepared from olefin 35 by an osmylation reaction with, for example, osmium tetroxide and oxidation of the resulting diol with, for example, Dess-Martin periodinane. This step is
5 followed by a saponification reaction with, for example, sodium hydroxide in methanol.


Syntheses of 39 and 40. These compounds are prepared according to Scheme 5 and experimentals previously described in Examples 1 and 2.

10

EXAMPLE 15

Sulfone-containing compounds of this invention (compounds 47 and 48) are prepared according to Scheme 6, below:

SCHEME 6. General Syntheses for the Sulfonamides



Synthesis of 42. This compound is prepared from bromoester 41 by treatment with magnesium in ether followed by addition of sulfur dioxide to the mixture.

5 **Synthesis of 43.** This compound is prepared from a mixture of sulfinate salt 42 and benzylcarbamate in water


by treatment with isobutyraldehyde in methanol. This step is followed by addition of formic acid with heating.


Synthesis of 43a. This compound is prepared by deprotection of the amino group of compound 43 with
5 hydrogen over palladium on carbon. This is followed by saponification of the ester with 5% TFA in dichloromethane, followed by treatment with Fmoc-OSU in dioxane-water.

Syntheses of 47 and 48. These compounds are
10 prepared according to Scheme 6 and experimentals previously described in Examples 1 and 2.

EXAMPLE 16

Sulfonyl-carbazate-containing compound of this invention (compounds 54 and 55) are prepared according to
15 Scheme 7, depicted below.

SCHEME 7. General Syntheses for the Sulfonyl Carbazates



Synthesis of 49. This compound is prepared from bromoester 41 by treatment with magnesium in ether, followed by addition of sulfur dioxide to the mixture and

subsequent treatment with sulfonyl chloride in dichloromethane at 0°C.


Synthesis of 50. This compound is prepared by coupling sulfinic acid 49 and Fmoc-isopropylcarbazate in the presence of DIEA in dichloromethane, followed by treatment with 5% TFA in dichloromethane.


Syntheses of 54 and 55. These compounds are prepared according to Scheme 7 and experimentals previously described in Examples 1 and 2.

10

EXAMPLE 17

Sulfonamide-containing compounds of this invention (compounds 62 and 63) are prepared according to Scheme 8, depicted below.


SCHEME 8. General Syntheses for the Sulfonamides




59

1) 25% pip / DMF
2) CapCO₂H / HBTU / HOBt / DIEA / NMP

1) 25% pip / DMF
2) Fmoc-Val / HBTU / HOBt / DIEA / NMP
3) 25% pip / DMF
4) Pyr-OH / HBTU / HOBt / DIEA / NMP




60




61

THF / H₂CO / AcOH /
0.1N HCl

THF / H₂CO / AcOH /
0.1N HCl



62



63

Synthesis of 56a. This compound is prepared from
5 ester 56 by treatment with lithium diisopropylamide in
THF at

-78°C followed by addition of sulfur dioxide to the mixture and subsequent treatment with sulfonyl chloride in dichloromethane at 0°C.

Synthesis of 57a. This compound is prepared by
5 coupling sulfonyl chloride 56a and anthranilate 57 in the presence of DIEA in dichloromethane.

Synthesis of 58. This compound is prepared by saponification of 57a with potassium hydroxide in ethanol followed by a Curtius rearrangement using, for example,
10 oxalyl chloride with sodium azide in acetonitrile. Protection of the amino functional group with Fmoc-OSU in dioxane-water, followed by hydrolysis of the *t*-butyl ester with 5% TFA in dichloromethane provide compound 58.

Syntheses of 62 and 63. These compounds could be
15 prepared according to Scheme 8 and experimentals previously described in Examples 1 and 2.

It will be readily apparent to those of skill in the art that other compounds of this invention may be synthesized using variations of Schemes 1 through 8
20 and/or the appropriate variants of the reagents depicted in those schemes and set forth in Examples 1 through 17.

Example 18

Inhibition of HCV NS3 Serine Protease

Insofar as compounds of formula (I) are able to
25 inhibit NS3 serine protease, they are of evident clinical utility for the treatment of viral diseases, including HCV. These tests are predictive of the compounds ability to inhibit HCV in vivo.

Peptides and Assays.

Peptides EDVVabuCSMSY (Abu designates - aminobutyric acid), DEMEECSQHLPYI, ECTTPCSGSWLRD and EDVV AbuC-p-nitroanilide was purchased from AnaSpec Inc. (San Jose, CA).

Peptide content of purified, lyophilized peptides and in-house peptides was determined by quantitative nitrogen analysis and the appropriate values were used in preparing stock peptide solutions (Galbreath). pKa determinations were determined by Robertson Microlit Laboratories, Inc. (Madison, NJ).

HPLC cleavage assays were performed using 25 nM to 3.0 μ M enzyme in 100 μ L volumes at 30 C containing 50 mM HEPES-KOH (pH 7.8), 100 mM NaCl, 20% glycerol, 5 mM DTT and the appropriate amount of substrate (in DMSO), with or without NS4A peptide, such that the final concentration of DMSO did not exceed 4%. Separate control experiments verified that this percentage of DMSO did not effect enzymatic activity. Cleavage reactions were quenched by the addition of an equal volume of a mixture of 10% TFA: acetonitrile (1:1) and activity was assessed on a reversed phase HPLC column (Rainin C18 Microsorb-MV, 5mm, 4.6 x 250mm; 0-50% acetonitrile, 0.1% TFA @ 3.33 min) using a Hewlett Packard 1050 instrument with auto-injection and diode array detection at 210 nm and 280 nm (where appropriate). Peptide elution fragments were collected and identified by mass spectrometry and N-terminal sequence analysis. Fragment identity and concentration was further verified by authentic, synthesized products. Initial rates of cleavage were determined at < 20% substrate conversion and catalytic parameters were determined assuming

Michaelis-Menten kinetics using the MultiFit program (Day Computing, Cambridge, MA).

Spectrophotometric assays were run in a 96-well microtiter plate at 30 C, using a SpectraMax 250 reader (Molecular Devices, Sunnyvale, CA) with kinetic capability. Cleavage of EDVV AbuC-p-nitroanilide (5A-pNA) substrate was performed with or without NS4A in the same buffer used for HPLC assays at 30 C, and pNA release was monitored at 405 nm. The extinction coefficient of p-nitroaniline is independent of pH at values of 5.5. and above [Tuppy, H., et al., Hoppe-Seyler's Z. Physiol. Chem., 329, pp. 278-288 (1962)]; Raybuck and Luong, unpublished observations). The percentage of DMSO did not exceed 4% in these assays.

Determination of the pH dependence of V_{max} , K_m and V_{max}/K_m was performed using a series of constant ionic strength buffers containing 50 mM MES, 25 mM Tris, 25 mM ethanolamine and 0.1 M NaCl [Morrison, J.F. and Stone, R.F., Biochemistry, 27, pp. 5499-5506 (1988)]. The inflection point for log V data was calculated by nonlinear least squares fit of the data to the equation.

$$\log v = \log[V_{max}/(1 + H/K_a)]$$

[Dixon, M. and Webb, E. C. Enzymes; Academic Press: New York; Vol., pp. 138-164 (1979)]. The inflection points for log (V/K) data were calculated by nonlinear least squares fit of the data to the equation:

$$\log v = \log[V_{max}/(1 + H/K_a + K_b/H)]$$

[Dixon, M. and Webb, E. C. Enzymes; Academic Press: New York; Vol., pp. 138-164 (1979)]. The program KineTic (BioKin Ltd) was used in both cases.

Kinetic constants for the rapid equilibrium ordered bisubstrate reaction were determined from rate

vs. [4A], [EDVV AbuC-pNA] data by non-linear least squares fitting to equation 1 [Morrison, J.F. Biochim. Biophys. Acta., 185, pp. 269-286 (1969)] as described in the text. K_{ii} and K_{is} values for peptidyl inhibitors were
 5 determined from rate vs. [inhibitor], [substrate] data and fitting to the equation for mixed inhibition:

$$\text{rate} = V_{\max}[S]/\{K_m(1+[I]/K_{is}) + [S] (1 + [I]/K_{ii})\}$$

The commercial program KinetAsyst (State College, PA) was used for both procedures. K_i values were calculated from
 10 rate vs. [inhibitor] plots by a nonlinear least squares fit of the data to the equation of Morrison for tight binding competitive inhibition [Morrison, J.F. Biochim. Biophys. Acta., 185, pp. 269-286 (1969)]. The KineTic program (BioKin Ltd) was used for this procedure.

15 The results are shown in Table 2. K_i values are expressed in μM . Category "A" indicates $< 1 \mu\text{M}$ inhibition; category "B" indicates 1-100 μM inhibition; category "C" indicates $> 100 \mu\text{M}$. The designation "ND" indicates that the compound was not tested.

20 Table 2. Enzyme inhibition data for compounds 1-9.

Compound	K_i	Compound	K_i	Compound	K_i
101	B	104	B	107	B
102	B	105	B	108	ND
103	B	106	B	109	ND


While we have hereinbefore presented a number of embodiments of this invention, it is apparent that my basic construction can be altered to provide other
 25 embodiments which utilize the methods of this invention. Therefore, it will be appreciated that the scope of this invention is to be defined by the claims appended hereto

rather than the specific embodiments which have been presented hereinbefore by way of example.

CLAIMS

What is claimed is:


1. A compound of the formula (I):



10

, wherein:

W is:



15

wherein:

m is 0 or 1;

20

each R¹ is hydroxy, alkoxy, or aryloxy, or each R¹ is an oxygen atom and together with the boron, to which they are each bound, form a 5-7 membered ring,

wherein the ring atoms are carbon, nitrogen, or oxygen;

each R^2 is independently hydrogen, alkyl, alkenyl, aryl, aralkyl, aralkenyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, heterocyclyl, heterocyclylalkyl, heterocyclylalkenyl, heteroaryl, or heteroaralkyl, or two R^2 groups, which are bound to the same nitrogen atom, form together with that nitrogen atom, a 5-7 membered monocyclic heterocyclic ring system; wherein any R^2 carbon atom is optionally substituted with J;

J is alkyl, aryl, aralkyl, alkoxy, aryloxy, aralkoxy, cycloalkyl, cycloalkoxy, heterocyclyl, heterocyclylalkyl, keto, hydroxy, amino, alkylamino, alkanoylamino, aroylamino, aralkanoylamino, carboxy, carboxyalkyl, carboxamidoalkyl, halo, cyano, nitro, formyl, acyl, sulfonyl, or sulfonamido and is optionally substituted with 1-3 J^1 groups; and

J^1 is alkyl, aryl, aralkyl, alkoxy, aryloxy, heterocyclyl, heterocyclylalkyl, keto, hydroxy, amino, alkanoylamino, aroylamino, carboxy, carboxyalkyl, carboxamidoalkyl, halo, cyano, nitro, formyl, sulfonyl, or sulfonamido;

L is alkyl, alkenyl, or alkynyl, wherein any hydrogen is optionally substituted with halogen, and wherein any hydrogen or halogen atom bound to any terminal carbon atom is optionally substituted with sulfhydryl or hydroxy;

each M is independently alkyl, cycloalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, or

heteroaralkyl, and is optionally substituted by 1 to 3 J groups, wherein any alkyl carbon atom may be replaced by a heteroatom;

R^{18} is a bond, $-N(R^{11})-$, or $-C(O)-$;

5 R^{11} is hydrogen or C_1-C_3 alkyl;

each R^{19} is independently H or R^{21} -aryl, or 2 adjacent R^{19} may be bound to one another to form a 5-7 membered aromatic ring; wherein any R^{19} is optionally substituted with 1 to 4 independently selected J' groups;

10 each R^{21} is independently C_1-C_3 -straight or branched alkyl, C_2-C_3 -straight or branched alkenyl, $O-(C_1-C_3)$ -straight or branched alkyl, or $O-(C_2-C_3)$ -straight or branched alkenyl;

n is 0 or 1;

15 the ring to which R^{18} and R^{19} are attached may be saturated, partially saturated, aromatic or fully unsaturated; and 1 to 3 carbon atoms that make up the ring to which R^{18} and R^{19} are attached are optionally replaced with a heteroatom which is independently

20 selected from O, S, $S(O)$, $S(O)_2$ or $N(R^{11})$;

A^2 is a bond or $-N(R^{11})-R^{17}(M)-R^{22}-$, wherein


R^{17} is $-CH-$ or $-N-$; and

R^{22} is $-C(O)-$ or $-S(O)_2-$;

V is a bond, $-CH(R^{11})-$, $-O-$, $-S-$, or $-N(R^{11})-$;

25 K is a bond, $-O-$, $-S-$, $-C(O)-$, $-S(O)-$, $-S(O)_2-$, or $-S(O)NR^{11}-$;

T is $-R^{12}$, $-alkyl-R^{12}$, $-alkenyl-R^{12}$, $-alkynyl-R^{12}$, $-OR^{12}$, $-N(R^{12})_2$, $-C(O)R^{12}$, $-C(=NO-alkyl)R^{12}$, or




each R^{12} is independently selected from
hydrogen, aryl, heteroaryl, cycloalkyl,
5 heterocyclyl, cycloalkylidenyl, or
heterocycloalkylidenyl, and is optionally
substituted with 1 to 3 J groups; or a first R^{12} and
a second R^{12} , together with the nitrogen to which
they are bound, form a mono- or bicyclic ring system
10 optionally substituted by 1 to 3 J groups;

R^{10} is alkyl, cycloalkyl, aryl, aralkyl,
heterocyclyl, heterocyclylalkyl, heteroaryl,
heteroaralkyl, carboxyalkyl, or carboxamidoalkyl,
and is optionally substituted with 1 to 3 hydrogens
15 J groups;

R^{15} is alkyl, cycloalkyl, aryl, aralkyl,
heterocyclyl, heterocyclylalkyl, heteroaryl,
heteroaralkyl, carboxyalkyl, or carboxamidoalkyl,
and is optionally substituted with 1 to 3 J groups;
20 and

R^{16} is hydrogen, alkyl, aryl, heteroaryl,
cycloalkyl, or heterocyclyl.

2. The compound according to claim 1, wherein
W is selected from:



5

3. The compound according to claim 2, wherein W is $-\text{C}(\text{O})\text{H}$.

4. The compound according to claim 1, wherein J is selected from alkyl, alkoxy, aryloxy, aryl, aralkyl, aralkoxy, halo, heteroaryl, cyano, amino, nitro, heterocyclyl, acyl, carboxy, carboxyalkyl, alkylamino, hydroxy, heterocyclylalkyl, aralkanoylamino, aroylamino, alkanoylamino, formyl or keto.

15

5. The compound according to claim 4, wherein J is selected from t-butyl, methyl, trifluoromethyl, hydroxy, methoxy, ethoxy, trifluoromethoxy, carboxy, phenyl, benzyl, phenoxy, benzyloxy, fluoro, chloro, bromo, isoxazolyl, pyridinyl, piperidinyl, carboxymethyl, carboxyethyl, dialkylamino, morpholinylmethyl,

phenylacetylamino, or acylamino.

6. The compound according to claim 1, wherein each J^1 is independently selected from alkoxy, alkyl, halo
5 or aryl.

7. The compound according to claim 6, wherein each J^1 is independently selected from C_{1-3} alkoxy, chloro, C_{1-3} alkyl, or phenyl.

10 8. The compound according to claim 1, wherein L is selected from trihalomethyl, sulfhydryl or alkyl substituted with trihalomethyl, sulfhydryl, or hydroxy.

9. The compound according to claim 8, wherein
15 L is $-CH_2CH_3$ or $-CH_2CF_3$.

10. The compound according to claim 1, wherein R^2 is selected from H, fluorine, trifluoromethyl, alkyl, aryl, aralkyl, heteroaralkyl, heterocyclyl, or
20 heterocyclylalkyl.

11. The compound according to claim 10, wherein R^2 is H.

25 12. The compound according to claim 1, wherein each M is independently selected from isopropyl, propyl, methyl, pyridylmethyl, benzyl, naphthylmethyl, phenyl, imidazolylmethyl, thiophenylmethyl, cyclohexylmethyl, phenethyl, benzylthiomethyl, or benzyloxyethyl.

30 13. The compound according to claim 12, wherein each M is isopropyl.

14. The compound according to claim 1, wherein
one R^{19} is R^{21} -aryl and the other two R^{19} are H, or two R^{19}
are bound together to form an aromatic ring and the other
5 R^{19} is H.

15. The compound according to claim 14,
wherein one R^{19} is -O-(C₁-C₃)-alkyl-aryl.

16. The compound according to claim 15,
wherein 14, wherein one R^{19} is -O-benzyl.

17. The compound according to claim 14,
15 wherein the two R^{19} that are bound together form a 6-
membered aromatic ring.

18. The compound according to claim 17,
wherein the two R^{19} that are bound together form phenyl.

19. The compound according to claim 1, wherein
20 R^{18} is -N(R^{11})-.

20. The compound according to claim 19,
25 wherein R^{18} is -N(H)- or -N(CH₃)-.

21. The compound according to claim 1, wherein
 A^2 is a bond or -N(R^{11})-C(M)-C(O)-.

22. The compound according to claim 21,
30 wherein A^2 is a bond or -N(H)-C(M)-C(O)-, wherein M is
isopropyl.


23. The compound according to claim 1, wherein
V is $-N(R^{11})-$.

5 24. The compound according to claim 23,
wherein V is $-NH-$.

25. The compound according to claim 1, wherein
K is $-C(O)-$ or $-S(O)_2-$.
10

26. The compound according to claim 25,
wherein K is $-C(O)-$.

27. The compound according to claim 1, wherein
15 T is selected from $-R^{12}$, $-alkyl-R^{12}$, $-alkenyl-R^{12}$, $-OR^{12}$,
 $-N(R^{12})_2$, $-C(=NO-alkyl)-R^{12}$, or



28. The compound according to claim 27,
20 wherein T is $-R^{12}$ or $-alkyl-R^{12}$.

29. The compound according to claim 1, wherein
 R^{12} is aryl or heteroaryl and is optionally substituted by
1-3 J groups.
25

30. The compound according to claim 29,
wherein R^{12} is naphthyl, pyrazinyl, or pyridyl, any of

which is optionally substituted with a hydroxy group.

31. The compound according to claim 1, wherein
R¹⁰ is alkyl substituted with carboxy.

5

32. The compound according to claim 1, wherein
R¹⁵ is alkyl substituted with carboxy.

33. The compound according to claim 1, wherein
10 the ring to which R¹⁸ and R¹⁹ are attached is aromatic.

34. A pharmaceutically acceptable composition
comprising:

a) a compound according to any one of claims 1-
15 33 in an amount effective to inhibit HCV NS3 protease;
and


b) a pharmaceutically suitable carrier.

35. The use of a compound according to any one
of claims 1-33 or a pharmaceutical composition according
20 to claim 34 in the manufacture of a medicament for
inhibiting serine protease activity in a patient.

36. The use according to claim 35, wherein the
serine protease is HCV NS3 protease.


37. The use of a compound according to any one
25 of claims 1-33 or a pharmaceutical composition according
to claim 34 in the manufacture of a medicament for
treating or preventing a hepatitis C viral infection in a
patient.

38. A process for preparing a compound of the formula (I):



, wherein:

W is:



wherein:

m is 0 or 1;

each R^1 is hydroxy, alkoxy, or aryloxy, or each R^1 is an oxygen atom and together with the boron, to which they are each bound, form a 5-7 membered ring,

wherein the ring atoms are carbon, nitrogen, or oxygen;

each R² is independently hydrogen, alkyl, alkenyl, aryl, aralkyl, aralkenyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, heterocyclyl, heterocyclylalkyl, heterocyclylalkenyl, heteroaryl, or heteroaralkyl, or two R² groups, which are bound to the same nitrogen atom, form together with that nitrogen atom, a 5-7 membered monocyclic heterocyclic ring system; wherein any R² carbon atom is optionally substituted with J;

J is alkyl, aryl, aralkyl, alkoxy, aryloxy, aralkoxy, cycloalkyl, cycloalkoxy, heterocyclyl, heterocyclylalkyl, keto, hydroxy, amino, alkylamino, alkanoylamino, aroylamino, aralkanoylamino, carboxy, carboxyalkyl, carboxamidoalkyl, halo, cyano, nitro, formyl, acyl, sulfonyl, or sulfonamido and is optionally substituted with 1-3 J¹ groups; and

J¹ is alkyl, aryl, aralkyl, alkoxy, aryloxy, heterocyclyl, heterocyclylalkyl, keto, hydroxy, amino, alkanoylamino, aroylamino, carboxy, carboxyalkyl, carboxamidoalkyl, halo, cyano, nitro, formyl, sulfonyl, or sulfonamido;

L is alkyl, alkenyl, or alkynyl, wherein any hydrogen is optionally substituted with halogen, and wherein any hydrogen or halogen atom bound to any terminal carbon atom is optionally substituted with sulfhydryl or hydroxy;

each M is independently alkyl, cycloalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, or

heteroaralkyl, and is optionally substituted by 1 to 3 J groups, wherein any alkyl carbon atom may be replaced by a heteroatom;

R^{18} is a bond, $-N(R^{11})-$, or $-C(O)-$;

5 R^{11} is hydrogen or C_1-C_3 alkyl;

each R^{19} is independently H or R^{21} -aryl, or 2 adjacent R^{19} may be bound to one another to form a 5-7 membered aromatic ring; wherein any R^{19} is optionally substituted with 1 to 4 independently selected J' groups;

10 each R^{21} is independently C_1-C_3 -straight or branched alkyl, C_2-C_3 -straight or branched alkenyl, $O-(C_1-C_3)$ -straight or branched alkyl, or $O-(C_2-C_3)$ -straight or branched alkenyl;

n is 0 or 1;

15 the ring to which R^{18} and R^{19} are attached may be saturated, partially saturated, aromatic or fully unsaturated; and 1 to 3 carbon atoms that make up the ring to which R^{18} and R^{19} are attached are optionally replaced with a heteroatom which is independently

20 selected from O, S, $S(O)$, $S(O)_2$ or $N(R^{11})$;

A^2 is a bond or $-N(R^{11})-R^{17}(M)-R^{22}-$, wherein


R^{17} is $-CH-$ or $-N-$; and

R^{22} is $-C(O)-$ or $-S(O)_2-$;

V is a bond, $-CH(R^{11})-$, $-O-$, $-S-$, or $-N(R^{11})-$;

25 K is a bond, $-O-$, $-S-$, $-C(O)-$, $-S(O)-$, $-S(O)_2-$, or $-S(O)NR^{11}-$;

T is $-R^{12}$, $-alkyl-R^{12}$, $-alkenyl-R^{12}$, $-alkynyl-R^{12}$, $-OR^{12}$, $-N(R^{12})_2$, $-C(O)R^{12}$, $-C(=NO-alkyl)R^{12}$, or



10

15

20


comprising the step of:

reacting a compound of formula (II):



, wherein LG is OH or an appropriate leaving group and the other substituents are as defined above;

5 with a compound of formula (III):



, wherein the NH₂ group is optionally protected and L and W are as defined above;

15 in the presence of a coupling reagent, provided that the compound of formula (II) or the compound of formula (III) is optionally bound to a resin.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/07149

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07C237/32 C07D241/24 C07D213/82 A61K31/495 A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	DE 196 48 793 A (BASF AG) 28 May 1998 (1998-05-28) claim 1	1
A	WO 96 33209 A (MERCK FROSSTCANADA INC.) 24 October 1996 (1996-10-24) claims 1,9	1,34

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

5 August 1999

Date of mailing of the international search report

11/08/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel.: (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Kapteyn, H

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/07149

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 1-38
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 99 07149

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1 - 38

Present claims 1 - 38 relate to an extremely large number of possible compounds. In fact, the claims contain so many options, variables, possible permutations that a lack of clarity and conciseness within the meaning of Article 6 PCT arises to such an extent as to render a meaningful search of the claims impossible. In addition the formula (I) claimed in claim 1 is not covered by the description. The central carbocycle of formula (1) is a saturated ring, but in all the examples there is an aromatic ring in this place. Consequently, the search has been carried out for those parts of the application which do appear to be clear and concise, namely examples 101 - 108 (Example 109 is not clear too)

From these examples the following values for the variables of formula (1) containing a central aromatic ring are derived:

A2-R18 = -NH-CH(isopropyl)CO-NH- or -NH-
K-V = -CO-NH-
L = unsubstituted ethyl or fluoro-substituted ethyl
M = isopropyl
R19A = H
R19B = H, -O-benzyl or -O-3,4-dichlorobenzyl
T = aryl or N-heteroaryl
W = -COH

The search has been executed in the scope of compounds covered by these variables and their preparation.

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

In ternational Application No

PCT/US 99/07149

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 19648793 A	28-05-1998	AU 5481498 A	22-06-1998
		WO 9823581 A	04-06-1998
		HR 970637 A	31-10-1998
<hr/>			
WO 9633209 A	24-10-1996	US 5798442 A	25-08-1998
		AU 5329096 A	07-11-1996
		CA 2218389 A	24-10-1996
<hr/>			